



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2024 – 05:10 pm BST

PDB ID : 4AKZ
Title : CRYSTAL STRUCTURE OF VIRB8 FROM BRUCELLA SUIS
Authors : Coincon, M.; Smith, M.A.; Sygusch, J.; Baron, C.
Deposited on : 2012-02-29
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	
1	B	138	
1	C	138	
1	D	138	
1	E	138	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 6291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYPE IV SECRETION SYSTEM PROTEIN VIRB8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	138	1094	688	181	222	3	0	0	0
1	B	134	1066	673	176	214	3	0	0	0
1	C	134	1066	673	176	214	3	0	0	0
1	D	133	1058	669	175	211	3	0	0	0
1	E	134	1066	673	176	214	3	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	218	Total 218	O 218	0	0
2	B	204	Total 204	O 204	0	0
2	C	152	Total 152	O 152	0	0
2	D	180	Total 180	O 180	0	0
2	E	187	Total 187	O 187	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8




- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8



- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8




- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8



- Molecule 1: TYPE IV SECRETION SYSTEM PROTEIN VIRB8





4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	198.44Å 198.44Å 103.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 2.25 19.84 – 2.24	Depositor EDS
% Data completeness (in resolution range)	97.6 (19.94-2.25) 97.7 (19.84-2.24)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.22 (at 2.23Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.190 , 0.217 0.223 , 0.241	Depositor DCC
R_{free} test set	3606 reflections (7.50%)	wwPDB-VP
Wilson B-factor (Å ²)	37.6	Xtrriage
Anisotropy	0.305	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 77.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6291	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	1/1119 (0.1%)	0.63	0/1522
1	B	0.61	0/1090	0.54	0/1481
1	C	0.60	0/1090	0.50	0/1481
1	D	0.52	0/1082	0.55	0/1470
1	E	0.49	0/1090	0.49	0/1481
All	All	0.60	1/5471 (0.0%)	0.54	0/7435

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	TYR	CD2-CE2	-5.95	1.30	1.39

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1094	0	1046	24	0
1	B	1066	0	1023	15	0
1	C	1066	0	1023	16	0
1	D	1058	0	1019	21	1
1	E	1066	0	1023	15	0
2	A	218	0	0	4	1
2	B	204	0	0	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	152	0	0	1	0
2	D	180	0	0	0	0
2	E	187	0	0	2	0
All	All	6291	0	5134	83	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:149:LYS:HD3	1:B:149:LYS:H	1.06	1.15
1:B:149:LYS:H	1:B:149:LYS:CD	1.71	1.04
1:B:149:LYS:CD	1:B:149:LYS:N	2.39	0.84
1:B:149:LYS:HD3	1:B:149:LYS:N	1.91	0.82
1:D:230:ARG:HD3	1:D:232:ASP:OD2	1.82	0.80
1:D:149:LYS:O	1:D:149:LYS:HG2	1.85	0.77
1:C:148:ASP:OD2	2:C:2069:HOH:O	2.05	0.74
1:D:97:SER:HB3	1:D:100:THR:CG2	2.20	0.70
1:A:188:ASN:HB2	2:A:2112:HOH:O	1.92	0.69
1:A:196:THR:HG22	2:A:2156:HOH:O	1.90	0.69
1:E:141:TYR:O	1:E:144:GLN:HG2	1.93	0.69
1:A:188:ASN:O	1:A:189:GLU:HB2	1.94	0.68
1:D:118:ASP:OD2	1:D:160:ARG:NH1	2.28	0.67
1:C:207:VAL:HG13	1:C:211:LEU:HD12	1.78	0.66
1:E:97:SER:O	1:E:101:VAL:HG23	1.98	0.63
1:A:219:THR:HB	2:A:2200:HOH:O	1.98	0.62
1:D:100:THR:O	1:D:104:LYS:HG3	1.99	0.62
1:A:141:TYR:O	1:A:144:GLN:HG2	2.01	0.61
1:E:115:GLU:HB3	1:E:182:LYS:HE3	1.84	0.60
1:A:139:GLN:NE2	1:D:175:ILE:HD11	2.17	0.60
1:A:209:PRO:HA	1:A:212:MET:HE2	1.85	0.59
1:A:121:THR:HG22	1:A:125:ASP:OD2	2.05	0.56
1:C:124:LYS:HD3	1:C:124:LYS:C	2.26	0.55
1:C:198:TRP:CZ3	1:C:233:PRO:HG3	2.41	0.55
1:A:196:THR:HG23	1:A:198:TRP:HE1	1.72	0.55
1:A:171:ASN:HD21	1:A:173:LYS:HB2	1.72	0.54
1:A:209:PRO:HA	1:A:212:MET:CE	2.37	0.54
1:C:121:THR:HG22	1:C:125:ASP:OD2	2.09	0.52
1:D:97:SER:HB3	1:D:100:THR:HG23	1.91	0.52
1:C:118:ASP:HB3	1:C:121:THR:HB	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLN:HG3	1:A:140:SER:N	2.25	0.52
1:E:196:THR:HG22	2:E:2100:HOH:O	2.09	0.52
1:A:131:MET:HG3	2:A:2201:HOH:O	2.08	0.52
1:A:121:THR:CG2	1:A:125:ASP:OD2	2.59	0.51
1:D:121:THR:HG23	1:D:125:ASP:OD2	2.11	0.51
1:A:144:GLN:HB3	1:D:230:ARG:NH2	2.26	0.51
1:A:189:GLU:HG2	1:A:190:THR:H	1.74	0.51
1:A:136:SER:O	1:A:139:GLN:HG3	2.11	0.50
1:C:98:TYR:C	1:C:98:TYR:CD1	2.83	0.50
1:D:179:ARG:NH1	1:D:234:GLU:OE1	2.44	0.50
1:B:207:VAL:O	1:B:208:ASN:C	2.48	0.50
1:B:196:THR:HG21	1:B:198:TRP:CZ2	2.48	0.48
1:C:155:TYR:HB3	1:C:159:VAL:HB	1.95	0.48
1:E:185:LYS:HE2	1:E:192:ASP:HA	1.96	0.48
1:B:196:THR:HG22	2:B:2163:HOH:O	2.13	0.48
1:A:189:GLU:HG2	1:A:190:THR:N	2.29	0.48
1:B:171:ASN:ND2	1:B:173:LYS:HB2	2.29	0.47
1:A:144:GLN:OE1	1:D:230:ARG:NH2	2.38	0.47
1:C:126:TYR:OH	1:C:142:ALA:HB1	2.16	0.46
1:C:212:MET:HG3	1:C:217:ARG:HG3	1.98	0.46
1:C:222:LEU:HD11	1:E:102:MET:HE1	1.97	0.46
1:C:166:VAL:HG22	1:D:166:VAL:HG22	1.98	0.46
1:C:196:THR:HB	1:C:198:TRP:HE1	1.81	0.46
1:E:97:SER:N	1:E:100:THR:HG1	2.14	0.45
1:B:98:TYR:CD1	1:B:98:TYR:C	2.90	0.45
1:B:196:THR:HB	1:B:198:TRP:HE1	1.82	0.45
1:B:98:TYR:HH	1:D:98:TYR:HE1	1.62	0.45
1:D:119:TRP:HB2	1:D:156:GLY:O	2.17	0.44
1:B:149:LYS:N	1:B:149:LYS:HD2	2.31	0.44
1:E:98:TYR:CD1	1:E:98:TYR:C	2.90	0.44
1:B:148:ASP:H	1:B:149:LYS:HD3	1.82	0.44
1:D:174:GLY:O	1:D:203:GLY:HA2	2.19	0.43
1:E:175:ILE:HG22	1:E:176:GLY:N	2.34	0.43
1:E:208:ASN:OD1	1:E:210:SER:OG	2.30	0.43
1:D:149:LYS:O	1:D:149:LYS:CG	2.60	0.43
1:E:155:TYR:N	1:E:155:TYR:CD1	2.85	0.43
1:C:186:ARG:O	1:C:187:THR:C	2.57	0.43
1:A:171:ASN:ND2	1:A:173:LYS:HB2	2.34	0.43
1:E:159:VAL:HG12	1:E:160:ARG:N	2.34	0.43
1:A:179:ARG:NH1	1:A:234:GLU:OE2	2.51	0.42
1:B:105:TYR:HB2	1:D:221:PRO:HG3	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:118:ASP:HB3	1:D:121:THR:HG22	2.01	0.42
1:A:139:GLN:HG3	1:A:140:SER:H	1.84	0.42
1:E:124:LYS:HG3	2:E:2021:HOH:O	2.18	0.42
1:C:168:ILE:HG12	1:C:178:VAL:HG22	2.02	0.42
1:E:104:LYS:HE2	1:E:168:ILE:O	2.20	0.42
1:D:182:LYS:O	1:D:195:THR:HA	2.19	0.41
1:B:123:GLN:O	1:B:127:GLU:HG3	2.20	0.41
1:E:209:PRO:HA	1:E:212:MET:HE3	2.02	0.41
1:D:102:MET:HG2	1:D:222:LEU:CD1	2.51	0.41
1:A:230:ARG:HH12	1:D:144:GLN:NE2	2.19	0.40
1:C:122:LEU:HD21	1:C:145:PHE:CE2	2.55	0.40
1:A:212:MET:HE1	1:A:223:GLY:HA2	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:131:MET:CG	2:A:2189:HOH:O[5_555]	2.09	0.11
2:B:2105:HOH:O	2:B:2105:HOH:O[5_554]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/138 (99%)	135 (99%)	0	1 (1%)	22	21
1	B	130/138 (94%)	130 (100%)	0	0	100	100
1	C	130/138 (94%)	127 (98%)	3 (2%)	0	100	100
1	D	129/138 (94%)	127 (98%)	2 (2%)	0	100	100
1	E	130/138 (94%)	126 (97%)	4 (3%)	0	100	100
All	All	655/690 (95%)	645 (98%)	9 (1%)	1 (0%)	47	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	189	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/121 (100%)	114 (94%)	7 (6%)	20	20
1	B	118/121 (98%)	115 (98%)	3 (2%)	47	56
1	C	118/121 (98%)	110 (93%)	8 (7%)	16	14
1	D	117/121 (97%)	111 (95%)	6 (5%)	24	25
1	E	118/121 (98%)	114 (97%)	4 (3%)	37	45
All	All	592/605 (98%)	564 (95%)	28 (5%)	26	29

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	121	THR
1	A	139	GLN
1	A	182	LYS
1	A	186	ARG
1	A	190	THR
1	A	194	GLU
1	A	196	THR
1	B	124	LYS
1	B	149	LYS
1	B	219	THR
1	C	98	TYR
1	C	108	SER
1	C	131	MET
1	C	139	GLN
1	C	144	GLN
1	C	148	ASP
1	C	196	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	207	VAL
1	D	100	THR
1	D	121	THR
1	D	157	SER
1	D	171	ASN
1	D	173	LYS
1	D	230	ARG
1	E	98	TYR
1	E	196	THR
1	E	215	SER
1	E	227	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/138 (100%)	0.45	12 (8%) 10 11	20, 32, 79, 111	0
1	B	134/138 (97%)	0.55	11 (8%) 11 12	23, 39, 78, 95	0
1	C	134/138 (97%)	0.67	16 (11%) 4 3	36, 54, 87, 110	0
1	D	133/138 (96%)	0.52	12 (9%) 9 10	22, 36, 70, 91	0
1	E	134/138 (97%)	0.89	24 (17%) 1 1	37, 52, 87, 106	0
All	All	673/690 (97%)	0.62	75 (11%) 5 5	20, 45, 84, 111	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	187	THR	8.2
1	C	97	SER	7.1
1	A	188	ASN	6.8
1	A	190	THR	6.8
1	E	192	ASP	6.0
1	C	148	ASP	5.7
1	A	191	GLY	5.6
1	E	149	LYS	5.4
1	C	147	GLY	5.2
1	B	173	LYS	5.2
1	C	187	THR	4.9
1	C	149	LYS	4.8
1	E	97	SER	4.8
1	B	97	SER	4.7
1	D	172	GLY	4.5
1	C	98	TYR	4.4
1	E	173	LYS	4.4
1	D	98	TYR	4.3
1	C	171	ASN	4.3
1	C	172	GLY	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	98	TYR	4.1
1	C	153	LYS	4.0
1	D	173	LYS	3.9
1	D	187	THR	3.8
1	E	148	ASP	3.8
1	A	189	GLU	3.7
1	E	230	ARG	3.7
1	E	186	ARG	3.6
1	E	194	GLU	3.5
1	E	193	GLY	3.5
1	D	148	ASP	3.5
1	E	171	ASN	3.5
1	E	150	ALA	3.4
1	E	233	PRO	3.4
1	E	98	TYR	3.3
1	E	147	GLY	3.2
1	A	187	THR	3.2
1	E	153	LYS	3.2
1	C	186	ARG	3.2
1	A	148	ASP	3.1
1	E	139	GLN	3.0
1	B	171	ASN	3.0
1	B	187	THR	3.0
1	D	171	ASN	3.0
1	C	194	GLU	2.9
1	E	154	GLN	2.9
1	A	97	SER	2.9
1	A	98	TYR	2.9
1	C	173	LYS	2.8
1	B	186	ARG	2.7
1	D	186	ARG	2.7
1	A	173	LYS	2.7
1	D	97	SER	2.6
1	B	192	ASP	2.6
1	C	120	TYR	2.6
1	A	192	ASP	2.6
1	E	99	ASP	2.6
1	C	146	GLN	2.6
1	D	193	GLY	2.5
1	B	178	VAL	2.5
1	E	119	TRP	2.5
1	D	149	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	171	ASN	2.4
1	D	156	GLY	2.4
1	C	119	TRP	2.3
1	B	149	LYS	2.3
1	B	194	GLU	2.2
1	D	194	GLU	2.2
1	E	146	GLN	2.2
1	C	99	ASP	2.2
1	E	178	VAL	2.1
1	E	124	LYS	2.1
1	B	198	TRP	2.0
1	E	145	PHE	2.0
1	A	153	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.