



# wwPDB X-ray Structure Validation Summary Report

Oct 23, 2021 – 09:29 AM EDT

PDB ID : 1CVU  
Title : CRYSTAL STRUCTURE OF ARACHIDONIC ACID BOUND TO THE CYCLOOXYGENASE ACTIVE SITE OF COX-2  
Authors : Kiefer, J.R.; Pawlitz, J.L.; Moreland, K.T.; Stegeman, R.A.; Gierse, J.K.; Stevens, A.M.; Goodwin, D.C.; Rowlinson, S.W.; Marnett, L.J.; Stallings, W.C.; Kurumbail, R.G.  
Deposited on : 1999-08-24  
Resolution : 2.40 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	552	 2% 68% 25% 7%
1	B	552	 2% 66% 27% 7%
2	F	9	 44% 44% 11%
3	C	2	 100%
3	E	2	 100%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	D	4	
4	G	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	2	-	-	-	X
3	NAG	E	2	-	-	-	X
4	MAN	D	4	X	-	-	X
4	MAN	G	3	-	-	-	X
4	MAN	G	4	X	-	-	X
5	NAG	A	681	-	-	-	X
5	NAG	B	2681	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 10100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROSTAGLANDIN H2 SYNTHASE-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	552	4469	2883	746	815	25	0	0	0
1	B	552	4469	2883	746	815	25	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	engineered mutation	UNP Q05769
A	333	LYS	ARG	engineered mutation	UNP Q05769
B	2310	GLN	ASN	engineered mutation	UNP Q05769
B	2333	LYS	ARG	engineered mutation	UNP Q05769

- Molecule 2 is a protein called PROTEIN (9-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	9	63	37	11	15	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0
3	E	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	4	50	28	2	20	0	0	0
4	G	4	50	28	2	20	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	B	1	14	8	1	5	0	0

- Molecule 6 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula:  $C_{14}H_{28}O_6$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C O	0	0
			20	14 6		
6	A	1	Total	C O	0	0
			20	14 6		
6	A	1	Total	C O	0	0
			20	14 6		
6	B	1	Total	C O	0	0
			20	14 6		
6	B	1	Total	C O	0	0
			20	14 6		

- Molecule 7 is ARACHIDONIC ACID (three-letter code: ACD) (formula:  $C_{20}H_{32}O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			22	20	2		
7	B	1	Total	C	O	0	0
			22	20	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	410	Total	O	0	0
			410	410		
8	B	361	Total	O	0	0
			361	361		





- Molecule 2: PROTEIN (9-MER)



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	181.04Å 133.96Å 124.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.40 34.95 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.1 (35.00-2.40) 96.2 (34.95-2.40)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.28 (at 2.39Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.204 , 0.235 0.195 , 0.224	Depositor DCC
$R_{free}$ test set	3887 reflections (6.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtrriage
Anisotropy	0.210	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 69.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10100	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ACD, BOG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	3/4595 (0.1%)	0.65	1/6229 (0.0%)
1	B	0.49	2/4595 (0.0%)	0.68	2/6229 (0.0%)
2	F	1.87	2/62 (3.2%)	1.70	2/82 (2.4%)
All	All	0.51	7/9252 (0.1%)	0.68	5/12540 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2603	SER	C-O	9.63	1.41	1.23
2	F	2603	SER	C-OXT	-7.12	1.09	1.23
1	A	203	GLN	CD-OE1	5.84	1.36	1.24
1	B	2170	GLU	CD-OE1	5.66	1.31	1.25
1	B	2170	GLU	CD-OE2	5.66	1.31	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	2603	SER	CA-C-O	11.83	144.95	120.10
1	B	2216	ARG	NE-CZ-NH1	-9.59	115.50	120.30
2	F	2603	SER	N-CA-C	6.91	129.66	111.00
1	B	2216	ARG	NE-CZ-NH2	5.32	122.96	120.30
1	A	148	TYR	N-CA-C	-5.18	97.01	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	409	TYR	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4469	0	4373	154	0
1	B	4469	0	4373	163	0
2	F	63	0	65	8	0
3	C	28	0	25	0	0
3	E	28	0	25	0	0
4	D	50	0	42	3	0
4	G	50	0	42	5	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
6	A	60	0	84	6	0
6	B	40	0	56	6	0
7	A	22	0	31	0	0
7	B	22	0	31	0	0
8	A	410	0	0	8	0
8	B	361	0	0	11	0
All	All	10100	0	9173	321	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 321 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2251:LYS:HD3	1:B:2310:GLN:HG3	1.49	0.92
1:A:251:LYS:HD3	1:A:310:GLN:HG3	1.51	0.89
1:B:2408:LEU:O	1:B:2409:TYR:HB2	1.74	0.87
1:B:2283:LEU:HD21	1:B:2415:LEU:HD12	1.55	0.85
1:A:543:GLN:OE1	1:A:543:GLN:HA	1.77	0.85

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/552 (100%)	522 (95%)	24 (4%)	4 (1%)	22	32
1	B	550/552 (100%)	516 (94%)	30 (6%)	4 (1%)	22	32
2	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	1107/1113 (100%)	1045 (94%)	54 (5%)	8 (1%)	22	32

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121	SER
1	A	122	TYR
1	B	2121	SER
1	B	2122	TYR
1	A	514	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/492 (100%)	445 (90%)	47 (10%)	8	12
1	B	492/492 (100%)	442 (90%)	50 (10%)	7	10
2	F	7/7 (100%)	6 (86%)	1 (14%)	3	4
All	All	991/991 (100%)	893 (90%)	98 (10%)	8	11

5 of 98 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2137	LYS
1	B	2298	LEU
1	B	2178	LEU
1	B	2252	LEU
1	B	2322	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	2454	GLN
2	F	2601	ASN
1	B	2034	ASN
1	B	2131	ASN
1	B	2242	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	C	1	1,3	14,14,15	0.80	0	17,19,21	0.70	0
3	NAG	C	2	3	14,14,15	0.92	0	17,19,21	0.93	0
4	NAG	D	1	1,4	14,14,15	0.39	0	17,19,21	0.65	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	D	2	4	14,14,15	2.04	2 (14%)	17,19,21	4.00	4 (23%)
4	MAN	D	3	4	11,11,12	1.35	2 (18%)	15,15,17	1.65	3 (20%)
4	MAN	D	4	4	11,11,12	2.27	3 (27%)	15,15,17	1.28	2 (13%)
3	NAG	E	1	1,3	14,14,15	0.83	0	17,19,21	0.66	0
3	NAG	E	2	3	14,14,15	0.87	0	17,19,21	0.91	0
4	NAG	G	1	1,4	14,14,15	0.66	0	17,19,21	0.89	0
4	NAG	G	2	4	14,14,15	1.30	2 (14%)	17,19,21	3.88	6 (35%)
4	MAN	G	3	4	11,11,12	1.15	1 (9%)	15,15,17	1.55	3 (20%)
4	MAN	G	4	4	11,11,12	2.30	3 (27%)	15,15,17	1.34	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	1/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	MAN	D	3	4	-	2/2/19/22	0/1/1/1
4	MAN	D	4	4	1/1/4/5	1/2/19/22	0/1/1/1
3	NAG	E	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	1/6/23/26	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	MAN	G	3	4	-	2/2/19/22	0/1/1/1
4	MAN	G	4	4	1/1/4/5	1/2/19/22	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	2	NAG	C8-C7	5.80	1.62	1.50
4	D	4	MAN	C2-C3	4.66	1.59	1.52
4	G	4	MAN	C2-C3	4.46	1.59	1.52
4	D	2	NAG	C1-C2	-4.07	1.46	1.52
4	G	4	MAN	C4-C5	3.95	1.61	1.53

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	G	2	NAG	C1-C2-N2	11.52	130.17	110.49
4	D	2	NAG	C1-C2-N2	10.93	129.16	110.49
4	D	2	NAG	C8-C7-N2	-10.64	98.08	116.10
4	G	2	NAG	C8-C7-N2	-8.98	100.90	116.10
4	G	2	NAG	O7-C7-C8	-4.87	113.02	122.06

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	4	MAN	C1
4	G	4	MAN	C1

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	2	NAG	C8-C7-N2-C2
4	D	2	NAG	O7-C7-N2-C2
4	G	2	NAG	C8-C7-N2-C2
4	G	2	NAG	O7-C7-N2-C2
4	G	3	MAN	C4-C5-C6-O6


There are no ring outliers.



5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	3	MAN	2	0
4	D	3	MAN	2	0
4	D	1	NAG	1	0
4	G	2	NAG	5	0
4	D	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
6	BOG	B	2702	-	20,20,20	0.68	0	25,25,25	0.61	0
5	NAG	A	681	1	14,14,15	0.87	1 (7%)	17,19,21	1.04	1 (5%)
6	BOG	A	703	-	20,20,20	0.92	0	25,25,25	0.67	0
6	BOG	B	2704	-	20,20,20	0.84	2 (10%)	25,25,25	0.66	0
5	NAG	B	2681	1	14,14,15	0.81	0	17,19,21	1.06	1 (5%)
7	ACD	A	701	-	18,21,21	0.66	0	17,21,21	0.53	0
7	ACD	B	2701	-	18,21,21	0.52	0	17,21,21	0.53	0
6	BOG	A	702	-	20,20,20	0.85	1 (5%)	25,25,25	0.67	0
6	BOG	A	704	-	20,20,20	0.93	2 (10%)	25,25,25	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BOG	B	2702	-	-	4/11/31/31	0/1/1/1
5	NAG	A	681	1	-	0/6/23/26	0/1/1/1
6	BOG	A	703	-	-	4/11/31/31	0/1/1/1
6	BOG	B	2704	-	-	5/11/31/31	0/1/1/1
5	NAG	B	2681	1	-	0/6/23/26	0/1/1/1
7	ACD	A	701	-	-	5/17/19/19	-
7	ACD	B	2701	-	-	2/17/19/19	-
6	BOG	A	702	-	-	2/11/31/31	0/1/1/1
6	BOG	A	704	-	-	6/11/31/31	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	704	BOG	O1-C1	2.39	1.44	1.40
6	A	704	BOG	O5-C1	2.34	1.47	1.41
6	B	2704	BOG	O5-C1	2.22	1.47	1.41
5	A	681	NAG	C1-C2	2.21	1.55	1.52
6	B	2704	BOG	O1-C1	2.19	1.43	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	2681	NAG	C8-C7-N2	-2.68	111.56	116.10
5	A	681	NAG	C8-C7-N2	2.17	119.77	116.10

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	2701	ACD	C12-C13-C14-C15
6	A	704	BOG	C4-C5-C6-O6
6	A	704	BOG	O5-C5-C6-O6
6	A	702	BOG	C4-C5-C6-O6
6	A	704	BOG	O5-C1-O1-C1'


There are no ring outliers.


5 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	2702	BOG	3	0
6	A	703	BOG	3	0
6	B	2704	BOG	3	0
6	A	702	BOG	1	0
6	A	704	BOG	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	552/552 (100%)	-0.36	11 (1%) 65 63	14, 26, 43, 56	0
1	B	552/552 (100%)	-0.28	13 (2%) 59 57	12, 27, 46, 61	0
2	F	9/9 (100%)	2.21	4 (44%) 0 0	52, 55, 66, 71	0
All	All	1113/1113 (100%)	-0.30	28 (2%) 57 55	12, 27, 46, 71	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2409	TYR	7.0
2	F	2603	SER	6.7
1	B	2583	GLN	4.0
1	B	2074	PHE	3.7
1	A	74	PHE	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MAN	G	4	11/12	0.38	0.78	78,80,82,82	0
4	MAN	G	3	11/12	0.54	0.42	66,67,72,76	0
4	MAN	D	3	11/12	0.58	0.40	60,62,65,67	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	E	2	14/15	0.63	0.70	68,71,74,75	0
3	NAG	C	2	14/15	0.65	0.48	64,67,70,72	0
4	MAN	D	4	11/12	0.66	0.43	70,71,72,72	0
3	NAG	E	1	14/15	0.67	0.35	52,54,58,64	0
4	NAG	G	2	14/15	0.78	0.36	49,55,57,62	0
4	NAG	D	2	14/15	0.80	0.36	46,52,55,57	0
3	NAG	C	1	14/15	0.86	0.23	44,50,53,58	0
4	NAG	G	1	14/15	0.94	0.15	27,29,34,41	0
4	NAG	D	1	14/15	0.95	0.13	28,30,33,39	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	BOG	A	704	20/20	0.57	0.35	42,63,65,67	0
5	NAG	B	2681	14/15	0.66	0.46	68,70,71,72	0
6	BOG	B	2704	20/20	0.77	0.25	32,61,64,65	0
5	NAG	A	681	14/15	0.79	0.42	58,61,64,66	0
6	BOG	A	702	20/20	0.81	0.30	38,50,54,54	0
6	BOG	B	2702	20/20	0.85	0.28	40,45,47,47	0
6	BOG	A	703	20/20	0.86	0.17	33,44,46,47	0
7	ACD	A	701	22/22	0.95	0.17	18,22,28,31	0
7	ACD	B	2701	22/22	0.95	0.15	19,25,29,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around BOG A 704:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around BOG B 2704:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around BOG A 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around BOG B 2702:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ACD A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.