



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 7, 2023 – 10:50 PM EDT

PDB ID : 6DWI
Title : Structure of the 4462 Antibody Fab fragment bound to a Staphylococcus aureus wall teichoic acid analog
Authors : Fong, R.; Lupardus, P.J.
Deposited on : 2018-06-26
Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	 2% 83% 8% 9%
1	C	233	 6% 75% 13% 12%
1	E	233	 4% 73% 14% 12%
1	G	233	 2% 82% 9% 9%
1	I	233	 2% 82% 9% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	233	 2% 79% 12% 9%
1	M	233	 % 78% 11% 11%
1	O	233	 78% 11% 9%
2	B	243	 % 75% 7% 17%
2	D	243	 4% 77% 9% 13%
2	F	243	 3% 74% 8% 18%
2	H	243	 % 77% 11% 13%
2	J	243	 2% 81% 7% 12%
2	L	243	 3% 75% 11% 13%
2	N	243	 % 76% 10% 14%
2	P	243	 76% 8% 16%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	H	302	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 26175 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4462 Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	211	Total 1619	C 1011	N 278	O 324	S 6	0	0	0
1	C	206	Total 1569	C 983	N 266	O 314	S 6	0	0	0
1	E	205	Total 1583	C 990	N 271	O 316	S 6	0	0	0
1	G	211	Total 1613	C 1008	N 275	O 324	S 6	0	0	0
1	I	211	Total 1613	C 1008	N 275	O 324	S 6	3	0	0
1	K	211	Total 1613	C 1008	N 275	O 324	S 6	0	0	0
1	M	207	Total 1582	C 990	N 272	O 314	S 6	0	0	0
1	O	211	Total 1619	C 1011	N 278	O 324	S 6	0	0	0

- Molecule 2 is a protein called 4462 Fab Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	202	Total 1523	C 962	N 259	O 295	S 7	4	0	0
2	D	212	Total 1579	C 993	N 270	O 309	S 7	4	0	0
2	F	200	Total 1508	C 952	N 256	O 293	S 7	4	1	0
2	H	212	Total 1586	C 998	N 270	O 311	S 7	0	1	0
2	J	214	Total 1601	C 1007	N 273	O 314	S 7	0	1	0
2	L	211	Total 1578	C 994	N 269	O 308	S 7	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	N	209	Total 1571	C 989	N 267	O 308	S 7	0	0	0
2	P	204	Total 1537	C 969	N 261	O 300	S 7	0	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0
3	B	1	Total 1	Cl 1	0	0
3	C	1	Total 1	Cl 1	0	0
3	F	1	Total 1	Cl 1	0	0
3	G	1	Total 1	Cl 1	0	0
3	O	1	Total 1	Cl 1	0	0

- Molecule 4 is 4-O-[2-acetamido-2-deoxy-beta-D-glucopyranosyl]-1-O-phosphono-D-ribitol (three-letter code: HD4) (formula: C₁₃H₂₆NO₁₃P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	D	1	Total 28	C 13	N 1	O 13	P 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
4	H	1	28	13	1	13	1	0	0
4	J	1	28	13	1	13	1	0	0
4	L	1	28	13	1	13	1	0	0

- Molecule 5 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	G	1	13	6	7	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	1	Total C O 6 3 3	0	0
6	J	1	Total C O 6 3 3	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	44	Total O 44 44	0	0
7	B	81	Total O 81 81	0	0
7	C	24	Total O 24 24	0	0
7	D	39	Total O 39 39	0	0
7	E	9	Total O 9 9	0	0
7	F	36	Total O 36 36	0	0
7	G	65	Total O 65 65	0	0
7	H	74	Total O 74 74	0	0
7	I	66	Total O 66 66	0	0
7	J	64	Total O 64 64	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	K	32	Total 32	O 32	0	0
7	L	33	Total 33	O 33	0	0
7	M	26	Total 26	O 26	0	0
7	N	42	Total 42	O 42	0	0
7	O	28	Total 28	O 28	0	0
7	P	75	Total 75	O 75	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4462 Fab Light Chain




- Molecule 1: 4462 Fab Light Chain



- Molecule 1: 4462 Fab Light Chain




- Molecule 1: 4462 Fab Light Chain







• Molecule 1: 4462 Fab Light Chain





• Molecule 1: 4462 Fab Light Chain




• Molecule 1: 4462 Fab Light Chain



• Molecule 1: 4462 Fab Light Chain




• Molecule 2: 4462 Fab Heavy chain







• Molecule 2: 4462 Fab Heavy chain




• Molecule 2: 4462 Fab Heavy chain





• Molecule 2: 4462 Fab Heavy chain




• Molecule 2: 4462 Fab Heavy chain



• Molecule 2: 4462 Fab Heavy chain



• Molecule 2: 4462 Fab Heavy chain



• Molecule 2: 4462 Fab Heavy chain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.57Å 114.10Å 128.30Å 90.00° 90.46° 90.00°	Depositor
Resolution (Å)	29.89 – 2.39 29.89 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.0 (29.89-2.39) 99.1 (29.89-2.39)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.39Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.203 , 0.249 0.207 , 0.259	Depositor DCC
R_{free} test set	6979 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.005 for l,k,-h 0.037 for h,-k,-l 0.014 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26175	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HD4, GOL, FLC, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/1654	0.72	0/2249
1	C	0.49	0/1601	0.73	0/2176
1	E	0.45	0/1616	0.69	0/2195
1	G	0.51	0/1648	0.74	0/2242
1	I	0.51	0/1648	0.74	0/2242
1	K	0.47	0/1648	0.70	0/2242
1	M	0.50	0/1615	0.73	0/2194
1	O	0.51	0/1654	0.74	0/2249
2	B	0.53	0/1558	0.75	0/2122
2	D	0.51	0/1615	0.73	0/2201
2	F	0.49	0/1545	0.73	0/2106
2	H	0.56	0/1625	0.78	0/2214
2	J	0.56	0/1640	0.77	0/2234
2	L	0.51	0/1614	0.75	0/2198
2	N	0.53	0/1607	0.78	0/2190
2	P	0.55	0/1572	0.75	0/2143
All	All	0.51	0/25860	0.74	0/35197

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1619	0	1577	5	0
1	C	1569	0	1516	15	0
1	E	1583	0	1540	18	0
1	G	1613	0	1566	9	0
1	I	1613	0	1566	9	0
1	K	1613	0	1566	11	0
1	M	1582	0	1531	14	0
1	O	1619	0	1577	20	0
2	B	1523	0	1489	10	0
2	D	1579	0	1532	12	0
2	F	1508	0	1463	9	0
2	H	1586	0	1549	13	0
2	J	1601	0	1564	5	0
2	L	1578	0	1544	9	0
2	N	1571	0	1539	5	0
2	P	1537	0	1498	8	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	F	1	0	0	1	0
3	G	1	0	0	0	0
3	O	1	0	0	0	0
4	D	28	0	0	0	0
4	H	28	0	0	1	0
4	J	28	0	0	0	0
4	L	28	0	0	0	0
5	G	13	0	5	0	0
6	H	6	0	8	0	0
6	J	6	0	8	0	0
7	A	44	0	0	0	0
7	B	81	0	0	0	0
7	C	24	0	0	1	0
7	D	39	0	0	0	0
7	E	9	0	0	0	0
7	F	36	0	0	0	0
7	G	65	0	0	1	0
7	H	74	0	0	0	0
7	I	66	0	0	0	0
7	J	64	0	0	0	0
7	K	32	0	0	1	0
7	L	33	0	0	0	0
7	M	26	0	0	0	0
7	N	42	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	O	28	0	0	0	0
7	P	75	0	0	0	0
All	All	26175	0	24638	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:47:TRP:HE1	2:B:50:THR:HG22	1.35	0.90
1:O:13:VAL:HG21	1:O:19:VAL:HG11	1.71	0.71
1:E:198:HIS:CG	1:E:199:GLN:H	2.09	0.70
1:O:29:VAL:HG11	1:O:90:GLN:HG2	1.72	0.70
2:L:11:LEU:HD21	2:L:150:PRO:HG3	1.74	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/233 (90%)	202 (97%)	7 (3%)	0	100	100
1	C	200/233 (86%)	189 (94%)	9 (4%)	2 (1%)	15	23
1	E	199/233 (85%)	188 (94%)	9 (4%)	2 (1%)	15	23
1	G	209/233 (90%)	203 (97%)	6 (3%)	0	100	100
1	I	209/233 (90%)	201 (96%)	8 (4%)	0	100	100
1	K	209/233 (90%)	200 (96%)	8 (4%)	1 (0%)	29	41
1	M	201/233 (86%)	191 (95%)	10 (5%)	0	100	100
1	O	209/233 (90%)	203 (97%)	6 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	196/243 (81%)	193 (98%)	2 (1%)	1 (0%)	29	41
2	D	208/243 (86%)	199 (96%)	8 (4%)	1 (0%)	29	41
2	F	195/243 (80%)	187 (96%)	8 (4%)	0	100	100
2	H	209/243 (86%)	202 (97%)	5 (2%)	2 (1%)	15	23
2	J	211/243 (87%)	206 (98%)	3 (1%)	2 (1%)	17	25
2	L	207/243 (85%)	196 (95%)	9 (4%)	2 (1%)	15	23
2	N	205/243 (84%)	199 (97%)	5 (2%)	1 (0%)	29	41
2	P	198/243 (82%)	192 (97%)	4 (2%)	2 (1%)	15	23
All	All	3274/3808 (86%)	3151 (96%)	107 (3%)	16 (0%)	29	41

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	138	ASN
2	D	99	ASN
2	H	99	ASN
2	H	129	PRO
2	J	99	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/199 (92%)	174 (96%)	8 (4%)	28	45
1	C	174/199 (87%)	162 (93%)	12 (7%)	15	25
1	E	179/199 (90%)	170 (95%)	9 (5%)	24	40
1	G	181/199 (91%)	173 (96%)	8 (4%)	28	45
1	I	181/199 (91%)	172 (95%)	9 (5%)	24	40
1	K	181/199 (91%)	175 (97%)	6 (3%)	38	57
1	M	175/199 (88%)	171 (98%)	4 (2%)	50	70
1	O	182/199 (92%)	177 (97%)	5 (3%)	44	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	172/207 (83%)	164 (95%)	8 (5%)	26	42
2	D	176/207 (85%)	168 (96%)	8 (4%)	27	44
2	F	169/207 (82%)	164 (97%)	5 (3%)	41	61
2	H	179/207 (86%)	172 (96%)	7 (4%)	32	50
2	J	181/207 (87%)	175 (97%)	6 (3%)	38	57
2	L	178/207 (86%)	165 (93%)	13 (7%)	14	22
2	N	179/207 (86%)	165 (92%)	14 (8%)	12	19
2	P	174/207 (84%)	164 (94%)	10 (6%)	20	33
All	All	2843/3248 (88%)	2711 (95%)	132 (5%)	27	43

5 of 132 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	N	200	ASN
1	O	78	LEU
2	P	189	SER
2	F	97	VAL
2	F	56	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HD4	H	301	-	28,28,28	0.96	1 (3%)	39,40,40	1.11	3 (7%)
4	HD4	L	301	-	28,28,28	0.97	1 (3%)	39,40,40	0.93	1 (2%)
6	GOL	H	302	-	5,5,5	0.11	0	5,5,5	0.20	0
5	FLC	G	302	-	12,12,12	1.05	0	17,17,17	1.31	2 (11%)
4	HD4	D	301	-	28,28,28	1.01	1 (3%)	39,40,40	1.06	0
4	HD4	J	301	-	28,28,28	1.10	1 (3%)	39,40,40	1.14	4 (10%)
6	GOL	J	302	-	5,5,5	0.17	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HD4	H	301	-	-	1/26/46/46	0/1/1/1
4	HD4	L	301	-	-	6/26/46/46	0/1/1/1
6	GOL	H	302	-	-	0/4/4/4	-
5	FLC	G	302	-	-	0/16/16/16	-
4	HD4	D	301	-	-	2/26/46/46	0/1/1/1
4	HD4	J	301	-	-	7/26/46/46	0/1/1/1
6	GOL	J	302	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	301	HD4	P27-O29	3.55	1.62	1.50
4	L	301	HD4	P27-O29	3.54	1.61	1.50
4	D	301	HD4	P27-O29	3.39	1.61	1.50
4	J	301	HD4	P27-O29	3.34	1.61	1.50

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	302	FLC	OB1-CBC-CB	-3.59	117.17	122.25
4	H	301	HD4	C1-O1-C17	2.93	119.86	115.33
4	J	301	HD4	C1-O1-C17	2.77	119.61	115.33
4	H	301	HD4	O5-C1-C2	-2.49	105.72	110.58
4	J	301	HD4	O1-C17-C21	2.40	112.02	107.47

There are no chirality outliers.

5 of 18 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	J	301	HD4	O1-C17-C19-O20
4	J	301	HD4	C21-C17-C19-O20
4	L	301	HD4	C17-C21-C23-C25
4	L	301	HD4	C17-C21-C23-O31
4	L	301	HD4	O32-C21-C23-C25


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	301	HD4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/233 (90%)	0.02	3 (1%) 75 73	28, 46, 64, 78	0
1	C	206/233 (88%)	0.27	14 (6%) 17 15	32, 53, 84, 97	0
1	E	205/233 (87%)	0.35	10 (4%) 29 28	40, 60, 81, 96	0
1	G	211/233 (90%)	-0.09	5 (2%) 59 57	19, 40, 66, 81	0
1	I	211/233 (90%)	-0.08	5 (2%) 59 57	23, 38, 67, 74	1 (0%)
1	K	211/233 (90%)	0.13	5 (2%) 59 57	32, 49, 69, 88	0
1	M	207/233 (88%)	-0.06	2 (0%) 82 80	25, 41, 66, 79	0
1	O	211/233 (90%)	0.01	0 100 100	26, 45, 67, 79	0
2	B	202/243 (83%)	-0.19	3 (1%) 73 72	20, 34, 63, 88	1 (0%)
2	D	212/243 (87%)	0.15	9 (4%) 36 35	29, 43, 68, 86	1 (0%)
2	F	200/243 (82%)	0.10	7 (3%) 44 43	32, 48, 69, 85	1 (0%)
2	H	212/243 (87%)	-0.17	3 (1%) 75 73	24, 35, 54, 91	0
2	J	214/243 (88%)	-0.12	6 (2%) 53 51	21, 35, 57, 76	0
2	L	211/243 (86%)	0.17	7 (3%) 46 45	28, 44, 81, 101	0
2	N	209/243 (86%)	-0.17	2 (0%) 82 80	27, 40, 63, 84	0
2	P	204/243 (83%)	-0.19	1 (0%) 91 89	22, 35, 59, 81	0
All	All	3337/3808 (87%)	0.01	82 (2%) 57 55	19, 43, 71, 101	4 (0%)

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	184	ALA	4.4
1	A	67	TYR	4.1
2	H	136	GLY	3.9
1	E	67	TYR	3.8
2	B	198	ILE	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	H	302	6/6	0.73	0.41	65,66,67,67	0
4	HD4	L	301	28/28	0.82	0.19	54,67,98,99	0
5	FLC	G	302	13/13	0.88	0.16	56,64,77,78	0
3	CL	O	301	1/1	0.89	0.09	43,43,43,43	0
4	HD4	J	301	28/28	0.92	0.14	32,44,68,69	0
4	HD4	D	301	28/28	0.94	0.11	38,46,69,71	0
6	GOL	J	302	6/6	0.94	0.14	42,46,49,50	0
4	HD4	H	301	28/28	0.96	0.10	26,38,52,53	0
3	CL	F	301	1/1	0.97	0.06	51,51,51,51	0
3	CL	B	301	1/1	0.98	0.08	38,38,38,38	0
3	CL	G	301	1/1	0.99	0.10	30,30,30,30	0
3	CL	C	301	1/1	0.99	0.15	35,35,35,35	0
3	CL	A	301	1/1	0.99	0.04	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HD4 L 301:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around HD4 J 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.