



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2024 – 06:57 AM EDT

PDB ID : 4IID  
Title : Crystal structure of beta-glucosidase 1 from *Aspergillus aculeatus* in complex with 1-deoxynojirimycin  
Authors : Suzuki, K.; Sumitani, J.; Kawaguchi, T.; Fushinobu, S.  
Deposited on : 2012-12-20  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	841	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>.</div> </div> </div>
1	B	841	<div> <div></div> <div>90%</div> <div>8%</div> <div>.</div> </div>
2	C	4	<div> <div></div> <div>100%</div> </div>
3	D	3	<div> <div></div> <div>67%</div> <div>33%</div> </div>
3	L	3	<div> <div></div> <div>100%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	N	3	 100%
4	E	2	 100%
4	G	2	 100%
5	F	10	 100%
5	M	10	 30% 70%
6	H	7	 29% 71%
6	O	7	 14% 57% 29%
7	I	7	 100%
8	J	7	 86% 14%
9	K	5	 100%
10	P	8	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
12	MRD	A	939	-	-	X	-

## 2 Entry composition [i](#)

There are 15 unique types of molecules in this entry. The entry contains 14782 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	834	Total	C	N	O	S	0	0	0
			6387	4031	1097	1241	18			
1	B	832	Total	C	N	O	S	0	0	0
			6375	4023	1095	1239	18			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	3	Total	C	N	O	0	0	0
			39	22	2	15			
3	L	3	Total	C	N	O	0	0	0
			39	22	2	15			
3	N	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	G	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	F	10	Total	C	N	O	0	0	0
			116	64	2	50			
5	M	10	Total	C	N	O	0	0	0
			116	64	2	50			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	H	7	Total	C	N	O	0	0	0
			83	46	2	35			
6	O	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	I	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	J	7	Total	C	N	O	0	0	0
			83	46	2	35			

- Molecule 9 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
9	K	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 10 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	P	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 11 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	N	O	0	0
			14	8	1	5		
11	A	1	Total	C	N	O	0	0
			14	8	1	5		
11	B	1	Total	C	N	O	0	0
			14	8	1	5		
11	B	1	Total	C	N	O	0	0
			14	8	1	5		
11	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 12 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	A	1	Total	C	O	0	0
			8	6	2		
12	A	1	Total	C	O	0	0
			8	6	2		
12	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 13 is 1-DEOXYNOJIRIMYCIN (three-letter code: NOJ) (formula:  $C_6H_{13}NO_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	A	1	Total	C	N	O	0	0
			11	6	1	4		


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	B	1	Total	C	N	O	0	0
			11	6	1	4		

- Molecule 14 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
14	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	450	Total	O	0	0
			450	450		
15	B	504	Total	O	0	0
			504	504		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-glucosidase 1




- Molecule 1: Beta-glucosidase 1



- Molecule 2: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  67% 33%


MAG1  
MAG2  
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%

MAG1  
MAG2  
BMA3

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:  100%


MAG1  
MAG2  
BMA3

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1  
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1  
MAG2

- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1  
MAG2  
BMA3  
MAN4  
MAN5  
MAN6  
MAN7  
MAN8  
MAN9  
MAN10

- Molecule 5: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  30% 70%





- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  29% 71%



- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:  14% 57% 29%




- Molecule 7: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%



- Molecule 8: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  86% 14%



- Molecule 9: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  100%



- Molecule 10:  $\alpha$ -D-mannopyranose-(1-2)- $\alpha$ -D-mannopyranose-(1-3)-[ $\alpha$ -D-mannopyranose-(1-6)] $\alpha$ -D-mannopyranose-(1-6)-[ $\alpha$ -D-mannopyranose-(1-3)] $\beta$ -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- $\beta$ -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- $\beta$ -D-glucopyranose

Chain P:

100%

NAG1  
NAG2  
BMA3  
MAN4  
MAN5  
MAN6  
MAN7  
MAN8

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.45Å 121.60Å 221.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.07 – 2.30 39.07 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (39.07-2.30) 99.5 (39.07-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.44 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.155 , 0.207 0.157 , 0.208	Depositor DCC
$R_{free}$ test set	4949 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 34.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14782	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NOJ, MPD, NAG, MRD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.94	1/6550 (0.0%)	0.93	11/8930 (0.1%)
1	B	1.01	5/6538 (0.1%)	0.94	13/8913 (0.1%)
All	All	0.97	6/13088 (0.0%)	0.94	24/17843 (0.1%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	541	CYS	CA-CB	6.87	1.69	1.53
1	A	541	CYS	CA-CB	6.63	1.68	1.53
1	B	820	ARG	CZ-NH2	6.09	1.41	1.33
1	B	511	TYR	CE1-CZ	5.67	1.46	1.38
1	B	541	CYS	CB-SG	5.49	1.91	1.82

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	521	ARG	NE-CZ-NH2	-16.07	112.27	120.30
1	A	521	ARG	NE-CZ-NH2	-15.81	112.39	120.30
1	A	521	ARG	NE-CZ-NH1	13.38	126.99	120.30
1	B	521	ARG	NE-CZ-NH1	12.77	126.69	120.30
1	A	70	LEU	CA-CB-CG	-8.84	94.98	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6387	0	6096	51	0
1	B	6375	0	6082	34	0
2	C	50	0	43	0	0
3	D	39	0	34	1	0
3	L	39	0	34	0	0
3	N	39	0	34	0	0
4	E	28	0	25	1	0
4	G	28	0	25	0	0
5	F	116	0	97	0	0
5	M	116	0	97	0	0
6	H	83	0	70	3	0
6	O	83	0	70	2	0
7	I	83	0	70	0	0
8	J	83	0	70	1	0
9	K	61	0	52	0	0
10	P	94	0	79	0	0
11	A	28	0	26	0	0
11	B	42	0	39	0	0
12	A	16	0	28	11	0
12	B	8	0	14	2	0
13	A	11	0	13	0	0
13	B	11	0	13	0	0
14	B	8	0	14	4	0
15	A	450	0	0	0	0
15	B	504	0	0	5	0
All	All	14782	0	13125	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 98 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:B:947:MPD:HM1	14:B:947:MPD:C5	1.66	1.16
14:B:947:MPD:H53	14:B:947:MPD:CM	1.79	1.11
1:A:63:THR:HG21	1:A:333:MET:CE	1.83	1.08
1:A:63:THR:HG21	1:A:333:MET:HE2	1.33	1.06
1:A:63:THR:CG2	1:A:333:MET:CE	2.39	1.01



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	830/841 (99%)	795 (96%)	35 (4%)	0	100	100
1	B	828/841 (98%)	799 (96%)	29 (4%)	0	100	100
All	All	1658/1682 (99%)	1594 (96%)	64 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	671/677 (99%)	663 (99%)	8 (1%)	71	84
1	B	670/677 (99%)	664 (99%)	6 (1%)	78	89
All	All	1341/1354 (99%)	1327 (99%)	14 (1%)	76	87

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	701	THR
1	B	37	GLN
1	B	853	HIS
1	B	700	LEU
1	B	757	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

78 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	2,1	14,14,15	1.02	1 (7%)	17,19,21	1.54	3 (17%)
2	NAG	C	2	2	14,14,15	0.95	1 (7%)	17,19,21	1.74	5 (29%)
2	BMA	C	3	2	11,11,12	1.26	1 (9%)	15,15,17	1.71	3 (20%)
2	MAN	C	4	2	11,11,12	1.02	0	15,15,17	2.09	3 (20%)
3	NAG	D	1	1,3	14,14,15	1.06	2 (14%)	17,19,21	1.49	3 (17%)
3	NAG	D	2	3	14,14,15	0.83	0	17,19,21	1.46	3 (17%)
3	BMA	D	3	3	11,11,12	1.08	0	15,15,17	2.67	7 (46%)
4	NAG	E	1	1,4	14,14,15	0.60	0	17,19,21	1.32	2 (11%)
4	NAG	E	2	4	14,14,15	0.60	0	17,19,21	1.60	3 (17%)
5	NAG	F	1	5,1	14,14,15	1.24	2 (14%)	17,19,21	0.72	0
5	MAN	F	10	5	11,11,12	1.12	2 (18%)	15,15,17	1.93	5 (33%)
5	NAG	F	2	5	14,14,15	0.78	0	17,19,21	1.29	2 (11%)
5	BMA	F	3	5	11,11,12	0.90	0	15,15,17	1.70	4 (26%)
5	MAN	F	4	5	11,11,12	0.63	0	15,15,17	1.68	2 (13%)
5	MAN	F	5	5	11,11,12	0.94	0	15,15,17	1.89	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MAN	F	6	5	11,11,12	0.92	1 (9%)	15,15,17	1.42	3 (20%)
5	MAN	F	7	5	11,11,12	0.64	0	15,15,17	1.43	2 (13%)
5	MAN	F	8	5	11,11,12	0.61	0	15,15,17	1.50	4 (26%)
5	MAN	F	9	5	11,11,12	0.75	0	15,15,17	1.11	2 (13%)
4	NAG	G	1	1,4	14,14,15	0.80	0	17,19,21	1.55	3 (17%)
4	NAG	G	2	4	14,14,15	0.88	1 (7%)	17,19,21	1.45	3 (17%)
6	NAG	H	1	1,6	14,14,15	0.69	0	17,19,21	0.83	0
6	NAG	H	2	6	14,14,15	0.93	1 (7%)	17,19,21	2.04	5 (29%)
6	BMA	H	3	6	11,11,12	0.83	0	15,15,17	1.07	0
6	MAN	H	4	6	11,11,12	0.79	0	15,15,17	0.99	0
6	MAN	H	5	6	11,11,12	0.84	0	15,15,17	1.76	5 (33%)
6	MAN	H	6	6	11,11,12	0.63	0	15,15,17	1.54	3 (20%)
6	MAN	H	7	6	11,11,12	0.69	0	15,15,17	1.46	2 (13%)
7	NAG	I	1	1,7	14,14,15	1.02	1 (7%)	17,19,21	1.94	4 (23%)
7	NAG	I	2	7	14,14,15	0.79	1 (7%)	17,19,21	1.33	2 (11%)
7	BMA	I	3	7	11,11,12	1.11	0	15,15,17	1.82	3 (20%)
7	MAN	I	4	7	11,11,12	0.92	0	15,15,17	1.35	2 (13%)
7	MAN	I	5	7	11,11,12	0.91	0	15,15,17	1.52	3 (20%)
7	MAN	I	6	7	11,11,12	0.57	0	15,15,17	1.98	5 (33%)
7	MAN	I	7	7	11,11,12	1.13	2 (18%)	15,15,17	3.45	7 (46%)
8	NAG	J	1	8,1	14,14,15	1.22	1 (7%)	17,19,21	1.54	4 (23%)
8	NAG	J	2	8	14,14,15	0.68	0	17,19,21	1.76	4 (23%)
8	BMA	J	3	8	11,11,12	0.94	0	15,15,17	1.67	2 (13%)
8	MAN	J	4	8	11,11,12	0.54	0	15,15,17	1.57	2 (13%)
8	MAN	J	5	8	11,11,12	0.89	0	15,15,17	1.27	1 (6%)
8	MAN	J	6	8	11,11,12	0.83	0	15,15,17	1.45	3 (20%)
8	MAN	J	7	8	11,11,12	1.01	0	15,15,17	2.30	8 (53%)
9	NAG	K	1	1,9	14,14,15	1.11	0	17,19,21	1.58	4 (23%)
9	NAG	K	2	9	14,14,15	0.84	0	17,19,21	1.56	4 (23%)
9	BMA	K	3	9	11,11,12	0.56	0	15,15,17	1.66	1 (6%)
9	MAN	K	4	9	11,11,12	1.07	1 (9%)	15,15,17	1.43	4 (26%)
9	MAN	K	5	9	11,11,12	0.93	0	15,15,17	1.64	3 (20%)
3	NAG	L	1	1,3	14,14,15	0.81	0	17,19,21	1.76	4 (23%)
3	NAG	L	2	3	14,14,15	0.73	0	17,19,21	1.69	4 (23%)
3	BMA	L	3	3	11,11,12	0.82	0	15,15,17	1.64	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	M	1	5,1	14,14,15	1.19	2 (14%)	17,19,21	1.84	6 (35%)
5	MAN	M	10	5	11,11,12	1.24	2 (18%)	15,15,17	1.92	6 (40%)
5	NAG	M	2	5	14,14,15	1.05	1 (7%)	17,19,21	1.54	3 (17%)
5	BMA	M	3	5	11,11,12	0.77	0	15,15,17	1.05	0
5	MAN	M	4	5	11,11,12	0.80	0	15,15,17	2.15	5 (33%)
5	MAN	M	5	5	11,11,12	0.58	0	15,15,17	2.35	5 (33%)
5	MAN	M	6	5	11,11,12	0.67	0	15,15,17	1.05	0
5	MAN	M	7	5	11,11,12	0.97	0	15,15,17	1.34	2 (13%)
5	MAN	M	8	5	11,11,12	1.41	2 (18%)	15,15,17	1.90	3 (20%)
5	MAN	M	9	5	11,11,12	0.85	0	15,15,17	0.73	0
3	NAG	N	1	1,3	14,14,15	0.76	0	17,19,21	1.36	1 (5%)
3	NAG	N	2	3	14,14,15	0.97	0	17,19,21	1.48	3 (17%)
3	BMA	N	3	3	11,11,12	0.89	0	15,15,17	2.14	4 (26%)
6	NAG	O	1	1,6	14,14,15	0.85	0	17,19,21	1.73	6 (35%)
6	NAG	O	2	6	14,14,15	1.16	1 (7%)	17,19,21	2.12	7 (41%)
6	BMA	O	3	6	11,11,12	0.59	0	15,15,17	2.95	4 (26%)
6	MAN	O	4	6	11,11,12	1.34	1 (9%)	15,15,17	2.01	4 (26%)
6	MAN	O	5	6	11,11,12	0.60	0	15,15,17	1.04	0
6	MAN	O	6	6	11,11,12	0.91	0	15,15,17	1.27	2 (13%)
6	MAN	O	7	6	11,11,12	1.02	1 (9%)	15,15,17	1.70	4 (26%)
10	NAG	P	1	1,10	14,14,15	0.71	0	17,19,21	1.96	3 (17%)
10	NAG	P	2	10	14,14,15	0.89	1 (7%)	17,19,21	1.44	1 (5%)
10	BMA	P	3	10	11,11,12	1.05	0	15,15,17	1.35	3 (20%)
10	MAN	P	4	10	11,11,12	0.70	0	15,15,17	1.22	1 (6%)
10	MAN	P	5	10	11,11,12	0.72	0	15,15,17	2.14	4 (26%)
10	MAN	P	6	10	11,11,12	0.71	0	15,15,17	1.74	5 (33%)
10	MAN	P	7	10	11,11,12	1.00	1 (9%)	15,15,17	1.59	4 (26%)
10	MAN	P	8	10	11,11,12	0.97	1 (9%)	15,15,17	1.63	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
3	BMA	D	3	3	-	2/2/19/22	0/1/1/1
4	NAG	E	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	2	4	-	3/6/23/26	0/1/1/1
5	NAG	F	1	5,1	-	0/6/23/26	0/1/1/1
5	MAN	F	10	5	-	2/2/19/22	0/1/1/1
5	NAG	F	2	5	-	0/6/23/26	0/1/1/1
5	BMA	F	3	5	-	0/2/19/22	0/1/1/1
5	MAN	F	4	5	-	0/2/19/22	0/1/1/1
5	MAN	F	5	5	-	1/2/19/22	0/1/1/1
5	MAN	F	6	5	-	0/2/19/22	0/1/1/1
5	MAN	F	7	5	-	0/2/19/22	0/1/1/1
5	MAN	F	8	5	-	0/2/19/22	0/1/1/1
5	MAN	F	9	5	-	0/2/19/22	0/1/1/1
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
6	NAG	H	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	H	2	6	-	0/6/23/26	0/1/1/1
6	BMA	H	3	6	-	0/2/19/22	0/1/1/1
6	MAN	H	4	6	-	0/2/19/22	0/1/1/1
6	MAN	H	5	6	-	0/2/19/22	0/1/1/1
6	MAN	H	6	6	-	0/2/19/22	0/1/1/1
6	MAN	H	7	6	-	2/2/19/22	0/1/1/1
7	NAG	I	1	1,7	-	1/6/23/26	0/1/1/1
7	NAG	I	2	7	-	0/6/23/26	0/1/1/1
7	BMA	I	3	7	-	0/2/19/22	0/1/1/1
7	MAN	I	4	7	-	2/2/19/22	0/1/1/1
7	MAN	I	5	7	-	0/2/19/22	0/1/1/1
7	MAN	I	6	7	-	2/2/19/22	0/1/1/1
7	MAN	I	7	7	-	2/2/19/22	0/1/1/1
8	NAG	J	1	8,1	-	0/6/23/26	0/1/1/1
8	NAG	J	2	8	-	2/6/23/26	0/1/1/1
8	BMA	J	3	8	-	0/2/19/22	0/1/1/1
8	MAN	J	4	8	-	0/2/19/22	0/1/1/1
8	MAN	J	5	8	-	1/2/19/22	0/1/1/1
8	MAN	J	6	8	-	2/2/19/22	0/1/1/1
8	MAN	J	7	8	-	0/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	K	1	1,9	-	0/6/23/26	0/1/1/1
9	NAG	K	2	9	-	0/6/23/26	0/1/1/1
9	BMA	K	3	9	-	0/2/19/22	0/1/1/1
9	MAN	K	4	9	-	2/2/19/22	0/1/1/1
9	MAN	K	5	9	-	1/2/19/22	0/1/1/1
3	NAG	L	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	L	2	3	-	2/6/23/26	0/1/1/1
3	BMA	L	3	3	-	2/2/19/22	0/1/1/1
5	NAG	M	1	5,1	-	0/6/23/26	0/1/1/1
5	MAN	M	10	5	-	0/2/19/22	0/1/1/1
5	NAG	M	2	5	-	0/6/23/26	0/1/1/1
5	BMA	M	3	5	-	0/2/19/22	0/1/1/1
5	MAN	M	4	5	-	0/2/19/22	0/1/1/1
5	MAN	M	5	5	-	0/2/19/22	0/1/1/1
5	MAN	M	6	5	-	0/2/19/22	0/1/1/1
5	MAN	M	7	5	-	0/2/19/22	0/1/1/1
5	MAN	M	8	5	-	0/2/19/22	0/1/1/1
5	MAN	M	9	5	-	0/2/19/22	0/1/1/1
3	NAG	N	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	N	2	3	-	2/6/23/26	0/1/1/1
3	BMA	N	3	3	-	1/2/19/22	0/1/1/1
6	NAG	O	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	O	2	6	-	0/6/23/26	0/1/1/1
6	BMA	O	3	6	-	0/2/19/22	0/1/1/1
6	MAN	O	4	6	-	0/2/19/22	0/1/1/1
6	MAN	O	5	6	-	1/2/19/22	0/1/1/1
6	MAN	O	6	6	-	1/2/19/22	0/1/1/1
6	MAN	O	7	6	-	1/2/19/22	0/1/1/1
10	NAG	P	1	1,10	-	1/6/23/26	0/1/1/1
10	NAG	P	2	10	-	0/6/23/26	0/1/1/1
10	BMA	P	3	10	-	0/2/19/22	0/1/1/1
10	MAN	P	4	10	-	1/2/19/22	0/1/1/1
10	MAN	P	5	10	-	0/2/19/22	0/1/1/1
10	MAN	P	6	10	-	2/2/19/22	0/1/1/1
10	MAN	P	7	10	-	0/2/19/22	0/1/1/1
10	MAN	P	8	10	-	2/2/19/22	0/1/1/1

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	O	4	MAN	O5-C1	-3.32	1.38	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	M	8	MAN	C2-C3	3.07	1.57	1.52
7	I	1	NAG	C2-N2	-3.05	1.41	1.46
5	F	1	NAG	O5-C1	-2.98	1.38	1.43
9	K	4	MAN	C2-C3	2.78	1.56	1.52

The worst 5 of 247 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	O	3	BMA	C6-C5-C4	-9.22	90.39	113.02
7	I	7	MAN	C1-O5-C5	8.15	123.11	112.19
10	P	5	MAN	C1-O5-C5	6.25	120.56	112.19
10	P	1	NAG	C1-O5-C5	6.00	120.23	112.19
7	I	7	MAN	C1-C2-C3	-5.80	101.20	109.64

There are no chirality outliers.

5 of 43 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
7	I	4	MAN	O5-C5-C6-O6
7	I	6	MAN	O5-C5-C6-O6
3	L	3	BMA	O5-C5-C6-O6
7	I	7	MAN	O5-C5-C6-O6
7	I	4	MAN	C4-C5-C6-O6

There are no ring outliers.



7 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	3	BMA	3	0
4	E	2	NAG	1	0
8	J	1	NAG	1	0
6	O	4	MAN	2	0
4	E	1	NAG	1	0
6	O	3	BMA	2	0
3	D	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





































## 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
11	NAG	B	908	1	14,14,15	0.79	0	17,19,21	1.25	1 (5%)
13	NOJ	A	940	-	11,11,11	0.48	0	13,15,15	1.19	2 (15%)
11	NAG	B	946	1	14,14,15	0.70	0	17,19,21	1.10	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
12	MRD	A	939	-	7,7,7	0.75	0	9,10,10	0.76	0
12	MRD	B	948	-	7,7,7	0.92	0	9,10,10	0.43	0
14	MPD	B	947	-	7,7,7	0.53	0	9,10,10	1.58	1 (11%)
11	NAG	A	905	1	14,14,15	0.94	0	17,19,21	1.62	3 (17%)
12	MRD	A	938	-	7,7,7	0.55	0	9,10,10	0.60	0
13	NOJ	B	949	-	11,11,11	1.47	2 (18%)	13,15,15	1.81	4 (30%)
11	NAG	B	945	1	14,14,15	0.59	0	17,19,21	1.42	2 (11%)
11	NAG	A	937	1	14,14,15	0.57	0	17,19,21	1.79	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	NAG	B	908	1	-	4/6/23/26	0/1/1/1
13	NOJ	A	940	-	-	0/2/19/19	0/1/1/1
11	NAG	B	946	1	-	2/6/23/26	0/1/1/1
12	MRD	A	939	-	-	4/5/5/5	-
12	MRD	B	948	-	-	2/5/5/5	-
14	MPD	B	947	-	-	0/5/5/5	-
11	NAG	A	905	1	-	0/6/23/26	0/1/1/1
12	MRD	A	938	-	-	1/5/5/5	-
13	NOJ	B	949	-	-	0/2/19/19	0/1/1/1
11	NAG	B	945	1	-	1/6/23/26	0/1/1/1
11	NAG	A	937	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	B	949	NOJ	C1-C2	3.20	1.55	1.52
13	B	949	NOJ	C2-C3	2.40	1.56	1.52

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	905	NAG	C1-C2-N2	4.11	116.91	110.43
11	B	945	NAG	O5-C1-C2	-3.66	105.64	111.29
11	A	937	NAG	O5-C5-C6	3.64	114.75	107.66
11	A	937	NAG	C3-C4-C5	-3.51	103.87	110.23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
13	B	949	NOJ	C1-N5-C5	3.40	117.18	109.71

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
12	A	939	MRD	C2-C3-C4-O4
12	A	939	MRD	C2-C3-C4-C5
12	B	948	MRD	C2-C3-C4-O4
12	B	948	MRD	C2-C3-C4-C5
11	B	946	NAG	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	A	939	MRD	9	0
12	B	948	MRD	2	0
14	B	947	MPD	4	0
12	A	938	MRD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	834/841 (99%)	-0.37	11 (1%) 77 81	11, 21, 36, 74	0
1	B	832/841 (98%)	-0.52	1 (0%) 95 97	11, 18, 32, 56	0
All	All	1666/1682 (99%)	-0.44	12 (0%) 87 91	11, 20, 34, 74	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	245	MET	3.0
1	A	144	ALA	2.8
1	A	675	VAL	2.7
1	A	277	VAL	2.6
1	A	189	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	MAN	K	4	11/12	0.68	0.25	71,78,86,88	0
9	MAN	K	5	11/12	0.74	0.28	56,66,70,73	0
6	MAN	O	7	11/12	0.76	0.29	71,76,79,85	0
10	MAN	P	7	11/12	0.78	0.23	45,61,66,70	0
3	BMA	L	3	11/12	0.80	0.28	57,75,80,91	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	MAN	H	7	11/12	0.81	0.34	66,68,75,77	0
10	MAN	P	8	11/12	0.82	0.22	62,72,77,79	0
7	MAN	I	7	11/12	0.83	0.23	51,57,67,69	0
3	BMA	N	3	11/12	0.85	0.26	56,65,69,69	0
2	MAN	C	4	11/12	0.87	0.17	42,50,53,58	0
4	NAG	G	2	14/15	0.88	0.21	49,56,64,65	0
7	MAN	I	6	11/12	0.89	0.19	50,56,59,62	0
10	MAN	P	6	11/12	0.90	0.23	51,58,63,67	0
2	BMA	C	3	11/12	0.91	0.09	32,39,46,47	0
5	MAN	M	10	11/12	0.91	0.12	32,38,45,48	0
6	BMA	H	3	11/12	0.91	0.19	32,35,43,53	0
6	MAN	O	4	11/12	0.92	0.17	30,35,41,41	0
3	BMA	D	3	11/12	0.92	0.25	38,43,52,56	0
6	NAG	O	2	14/15	0.92	0.18	23,28,38,38	0
6	BMA	O	3	11/12	0.92	0.17	34,39,47,54	0
9	BMA	K	3	11/12	0.92	0.17	47,53,62,78	0
6	NAG	H	2	14/15	0.93	0.15	18,26,37,39	0
10	NAG	P	2	14/15	0.93	0.13	25,35,42,44	0
10	BMA	P	3	11/12	0.93	0.15	37,44,55,66	0
8	MAN	J	7	11/12	0.93	0.12	34,38,42,44	0
7	BMA	I	3	11/12	0.93	0.13	38,43,49,53	0
5	MAN	M	6	11/12	0.93	0.13	29,37,42,43	0
5	MAN	F	6	11/12	0.94	0.17	33,38,43,43	0
6	MAN	H	4	11/12	0.94	0.17	30,36,41,46	0
5	MAN	F	10	11/12	0.94	0.17	35,45,55,58	0
3	NAG	N	2	14/15	0.94	0.18	39,42,50,57	0
8	MAN	J	5	11/12	0.94	0.19	33,37,46,61	0
3	NAG	L	2	14/15	0.94	0.14	20,32,45,51	0
3	NAG	D	2	14/15	0.94	0.12	24,31,38,39	0
4	NAG	G	1	14/15	0.95	0.10	29,36,44,47	0
8	BMA	J	3	11/12	0.95	0.09	26,28,32,32	0
7	NAG	I	2	14/15	0.95	0.13	26,32,42,45	0
6	NAG	H	1	14/15	0.95	0.14	18,26,39,42	0
7	MAN	I	4	11/12	0.95	0.10	27,34,41,52	0
6	NAG	O	1	14/15	0.95	0.11	16,24,41,41	0
6	MAN	O	6	11/12	0.96	0.13	26,31,37,39	0
8	MAN	J	6	11/12	0.96	0.14	28,31,36,40	0
5	MAN	F	7	11/12	0.96	0.09	23,25,27,27	0
9	NAG	K	2	14/15	0.96	0.13	27,29,35,42	0
4	NAG	E	2	14/15	0.96	0.13	32,37,42,47	0
5	NAG	M	2	14/15	0.96	0.09	18,20,27,30	0
2	NAG	C	2	14/15	0.96	0.09	27,31,36,39	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
10	NAG	P	1	14/15	0.96	0.08	19,23,26,27	0
3	NAG	D	1	14/15	0.96	0.08	19,25,30,34	0
5	MAN	F	5	11/12	0.96	0.13	24,28,35,39	0
10	MAN	P	4	11/12	0.96	0.16	27,35,47,52	0
10	MAN	P	5	11/12	0.96	0.13	26,33,38,48	0
8	NAG	J	1	14/15	0.96	0.09	14,17,20,21	0
4	NAG	E	1	14/15	0.96	0.08	18,24,27,29	0
8	MAN	J	4	11/12	0.96	0.16	28,34,39,41	0
7	NAG	I	1	14/15	0.97	0.08	21,24,26,31	0
9	NAG	K	1	14/15	0.97	0.09	21,24,27,28	0
5	NAG	F	2	14/15	0.97	0.10	26,27,31,31	0
5	NAG	M	1	14/15	0.97	0.08	17,20,23,25	0
6	MAN	H	6	11/12	0.97	0.17	29,31,40,49	0
7	MAN	I	5	11/12	0.97	0.11	28,35,45,49	0
2	NAG	C	1	14/15	0.97	0.09	19,25,26,27	0
5	BMA	M	3	11/12	0.97	0.07	20,22,25,26	0
3	NAG	N	1	14/15	0.97	0.08	26,33,46,54	0
8	NAG	J	2	14/15	0.97	0.08	16,21,25,31	0
5	MAN	M	8	11/12	0.97	0.08	18,20,20,25	0
5	NAG	F	1	14/15	0.97	0.12	24,28,31,31	0
5	MAN	F	8	11/12	0.97	0.13	21,22,26,27	0
5	MAN	F	9	11/12	0.97	0.13	30,34,38,41	0
5	BMA	F	3	11/12	0.98	0.09	24,26,29,31	0
5	MAN	M	9	11/12	0.98	0.07	19,22,24,31	0
5	MAN	F	4	11/12	0.98	0.08	17,20,23,23	0
5	MAN	M	4	11/12	0.98	0.08	17,19,20,21	0
5	MAN	M	5	11/12	0.98	0.08	21,23,30,32	0
3	NAG	L	1	14/15	0.98	0.07	15,19,22,23	0
6	MAN	O	5	11/12	0.98	0.14	27,30,34,40	0
5	MAN	M	7	11/12	0.98	0.07	15,17,20,24	0
6	MAN	H	5	11/12	0.98	0.15	25,27,28,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around Chain D:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain L:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain N:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain G:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain F:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain M:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain H:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain O:**



$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around Chain I:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around Chain J:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
11	NAG	A	905	14/15	0.83	0.32	49,63,72,73	0
11	NAG	B	908	14/15	0.90	0.28	44,58,67,69	0
11	NAG	B	945	14/15	0.91	0.24	45,60,67,70	0
12	MRD	A	939	8/8	0.93	0.14	24,28,37,37	0
14	MPD	B	947	8/8	0.93	0.14	26,32,40,44	0
12	MRD	B	948	8/8	0.95	0.16	38,45,48,52	0
12	MRD	A	938	8/8	0.96	0.11	36,38,40,41	0
11	NAG	A	937	14/15	0.96	0.14	39,45,50,52	0
11	NAG	B	946	14/15	0.97	0.22	29,35,40,43	0
13	NOJ	B	949	11/11	0.98	0.17	14,15,17,18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
13	NOJ	A	940	11/11	0.99	0.23	17,21,22,24	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.