



Full wwPDB X-ray Structure Validation Report i

Jun 16, 2024 – 07:52 PM EDT

PDB ID : 5J7X
Title : Baeyer-Villiger monooxygenase BVMOAFL838 from Aspergillus flavus
Authors : Ferroni, F.M.; Tolmie, C.; Smit, M.S.; Opperman, D.J.
Deposited on : 2016-04-07
Resolution : 1.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	549	1% 79%	15% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TAM	A	603	-	X	-	-

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dimethylaniline monooxygenase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	527	4229	2712	713	790	14	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	544	HIS	-	expression tag	UNP B8N653
A	545	HIS	-	expression tag	UNP B8N653
A	546	HIS	-	expression tag	UNP B8N653
A	547	HIS	-	expression tag	UNP B8N653
A	548	HIS	-	expression tag	UNP B8N653
A	549	HIS	-	expression tag	UNP B8N653

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0

- Molecule 4 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: C₇H₁₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	7	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	361	Total O		0	0
			361	361		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dimethylaniline monooxygenase, putative



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	109.67 Å 177.51 Å 76.37 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.40 – 1.90 32.42 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (32.40-1.90) 99.5 (32.42-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.76 (at 1.89 Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R , R_{free}	0.157 , 0.204 0.168 , 0.211	Depositor DCC
R_{free} test set	2981 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.6	EDS
L-test for twinning ²	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4659	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TAM, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	1.62	40/4346 (0.9%)	1.39	44/5906 (0.7%)

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	GLU	CD-OE1	13.46	1.40	1.25
1	A	486	GLU	CD-OE2	10.13	1.36	1.25
1	A	278	TYR	CG-CD2	-9.78	1.26	1.39
1	A	278	TYR	CZ-OH	8.82	1.52	1.37
1	A	75	GLU	CD-OE2	8.77	1.35	1.25
1	A	389	GLY	CA-C	8.54	1.65	1.51
1	A	162	SER	CB-OG	8.50	1.53	1.42
1	A	141	TYR	CE1-CZ	-8.37	1.27	1.38
1	A	439	GLN	C-O	7.17	1.36	1.23
1	A	46	SER	CB-OG	7.16	1.51	1.42
1	A	331	TYR	CZ-OH	6.91	1.49	1.37
1	A	480	GLU	CD-OE2	-6.69	1.18	1.25
1	A	526	ARG	CZ-NH1	6.63	1.41	1.33
1	A	45	GLU	CB-CG	6.55	1.64	1.52
1	A	61	ARG	CZ-NH2	-6.42	1.24	1.33
1	A	45	GLU	CG-CD	6.39	1.61	1.51
1	A	436	TYR	CG-CD2	6.34	1.47	1.39
1	A	75	GLU	CD-OE1	6.24	1.32	1.25
1	A	147	GLY	CA-C	6.23	1.61	1.51
1	A	412	GLU	CD-OE2	6.18	1.32	1.25
1	A	114	ARG	CZ-NH2	-6.13	1.25	1.33
1	A	69	TYR	CG-CD2	-6.00	1.31	1.39
1	A	132	THR	N-CA	-5.80	1.34	1.46
1	A	94	SER	CA-CB	5.76	1.61	1.52
1	A	413	ARG	CZ-NH2	-5.62	1.25	1.33
1	A	360	SER	CA-CB	5.58	1.61	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	GLU	CG-CD	5.51	1.60	1.51
1	A	269	GLU	CB-CG	-5.50	1.41	1.52
1	A	32	ARG	CZ-NH1	5.50	1.40	1.33
1	A	341	GLU	CB-CG	5.35	1.62	1.52
1	A	486	GLU	CG-CD	5.32	1.59	1.51
1	A	405	VAL	C-O	-5.30	1.13	1.23
1	A	171	ALA	N-CA	5.29	1.56	1.46
1	A	22	PHE	CG-CD2	5.29	1.46	1.38
1	A	170	SER	C-O	-5.25	1.13	1.23
1	A	502	TRP	CD2-CE2	5.15	1.47	1.41
1	A	91	GLU	CG-CD	5.14	1.59	1.51
1	A	69	TYR	CE2-CZ	5.11	1.45	1.38
1	A	22	PHE	CE1-CZ	5.09	1.47	1.37
1	A	275	GLU	CD-OE2	5.07	1.31	1.25

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	504	MET	CG-SD-CE	-17.11	72.82	100.20
1	A	526	ARG	NE-CZ-NH2	-14.81	112.90	120.30
1	A	317	ASP	CB-CG-OD1	11.58	128.73	118.30
1	A	526	ARG	NE-CZ-NH1	10.53	125.56	120.30
1	A	30	LYS	CD-CE-NZ	8.65	131.59	111.70
1	A	456	ASP	CB-CG-OD1	8.56	126.01	118.30
1	A	382	ASP	CB-CG-OD1	-7.86	111.23	118.30
1	A	328	GLU	OE1-CD-OE2	-7.75	114.00	123.30
1	A	480	GLU	OE1-CD-OE2	-7.50	114.30	123.30
1	A	411	ASP	CB-CG-OD1	7.47	125.03	118.30
1	A	284	ARG	NE-CZ-NH1	7.34	123.97	120.30
1	A	14	ASP	CB-CG-OD1	-7.29	111.74	118.30
1	A	411	ASP	CB-CG-OD2	-7.12	111.89	118.30
1	A	114	ARG	NE-CZ-NH1	6.73	123.66	120.30
1	A	500	ASP	CB-CG-OD1	6.44	124.10	118.30
1	A	297	LEU	CA-CB-CG	6.33	129.86	115.30
1	A	536	ASP	CB-CG-OD1	6.26	123.93	118.30
1	A	412	GLU	OE1-CD-OE2	6.23	130.78	123.30
1	A	75	GLU	OE1-CD-OE2	6.22	130.77	123.30
1	A	292	ASP	CB-CG-OD2	-6.17	112.75	118.30
1	A	317	ASP	CB-CG-OD2	-6.15	112.77	118.30
1	A	101	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	A	126	ARG	NE-CZ-NH1	6.09	123.34	120.30
1	A	269	GLU	CG-CD-OE2	-6.09	106.13	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	337	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	160	LEU	CB-CA-C	-5.94	98.91	110.20
1	A	489	MET	CG-SD-CE	5.90	109.63	100.20
1	A	474	THR	CA-CB-CG2	-5.88	104.17	112.40
1	A	32	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	175	GLU	N-CA-C	-5.75	95.47	111.00
1	A	75	GLU	CG-CD-OE1	-5.69	106.92	118.30
1	A	467	MET	CG-SD-CE	5.68	109.29	100.20
1	A	343	ASP	CB-CG-OD1	5.58	123.32	118.30
1	A	61	ARG	NE-CZ-NH1	5.55	123.08	120.30
1	A	31	LEU	CB-CG-CD2	5.54	120.43	111.00
1	A	438	LEU	CA-CB-CG	5.53	128.02	115.30
1	A	321	ARG	N-CA-CB	-5.41	100.86	110.60
1	A	47	ASP	CB-CG-OD1	5.34	123.11	118.30
1	A	468	GLU	OE1-CD-OE2	-5.31	116.92	123.30
1	A	489	MET	CA-CB-CG	-5.27	104.34	113.30
1	A	45	GLU	CB-CA-C	5.26	120.91	110.40
1	A	215	THR	CB-CA-C	-5.16	97.68	111.60
1	A	121	ASP	CB-CG-OD2	-5.09	113.72	118.30
1	A	337	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4229	0	4089	27	0
2	A	53	0	31	0	0
3	A	5	0	0	0	0
4	A	11	0	15	2	0
5	A	361	0	0	1	2
All	All	4659	0	4135	27	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:GLN:HA	1:A:78:GLN:OE1	1.76	0.82
1:A:393:VAL:HG11	1:A:438:LEU:HB3	1.65	0.77
1:A:214:ARG:HB3	4:A:603:TAM:H62	1.76	0.67
1:A:500:ASP:O	1:A:504:MET:HG2	1.97	0.65
1:A:175:GLU:O	1:A:176:ALA:CB	2.48	0.61
1:A:78:GLN:OE1	1:A:78:GLN:CA	2.50	0.60
1:A:414:TRP:CH2	1:A:438:LEU:HD21	2.42	0.55
1:A:79:SER:HB2	5:A:877:HOH:O	2.09	0.53
1:A:175:GLU:O	1:A:176:ALA:HB3	2.09	0.52
1:A:185:LYS:HE3	1:A:207:GLU:OE2	2.11	0.50
1:A:75:GLU:H	1:A:75:GLU:CD	2.15	0.50
1:A:367:THR:HB	1:A:368:PRO:CD	2.43	0.49
1:A:457:TRP:CD1	1:A:529:CYS:HB3	2.48	0.48
1:A:160:LEU:HD22	1:A:167:ILE:CD1	2.45	0.47
1:A:393:VAL:CG1	1:A:438:LEU:HB3	2.41	0.45
1:A:461:LEU:C	1:A:461:LEU:HD23	2.37	0.45
1:A:386:VAL:HG12	1:A:388:THR:HG23	1.99	0.45
1:A:453:LEU:C	1:A:453:LEU:HD23	2.37	0.45
1:A:74:PRO:O	1:A:78:GLN:HG2	2.18	0.43
1:A:367:THR:HB	1:A:368:PRO:HD2	1.99	0.43
1:A:245:GLU:HG3	1:A:246:CYS:N	2.34	0.42
1:A:352:ASN:OD1	1:A:352:ASN:N	2.42	0.42
1:A:17:ILE:HB	1:A:41:ILE:HG12	2.00	0.42
1:A:394:THR:O	1:A:398:LEU:HG	2.20	0.41
1:A:148:PHE:CE2	1:A:393:VAL:HB	2.56	0.41
1:A:214:ARG:HE	4:A:603:TAM:H52	1.86	0.41
1:A:166:THR:HG21	1:A:168:TYR:CZ	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:716:HOH:O	5:A:716:HOH:O[2_775]	0.94	1.26
5:A:1037:HOH:O	5:A:1037:HOH:O[3_756]	1.23	0.97

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	524/549 (95%)	509 (97%)	13 (2%)	2 (0%)	34 24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	ALA
1	A	442	THR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	447/464 (96%)	442 (99%)	5 (1%)	73 73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	175	GLU
1	A	205	GLU
1	A	355	ILE
1	A	438	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	257	GLN
1	A	406	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	602	-	4,4,4	1.24	1 (25%)	6,6,6	0.99	0
2	FAD	A	601	-	53,58,58	1.82	13 (24%)	68,89,89	1.74	18 (26%)
4	TAM	A	603	-	7,10,10	1.49	1 (14%)	9,12,12	6.32	7 (77%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	5/30/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TAM	A	603	-	-	7/12/12/12	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C1'-C2'	5.82	1.60	1.52
2	A	601	FAD	C9A-C5X	4.80	1.49	1.41
2	A	601	FAD	C5'-C4'	3.93	1.57	1.51
2	A	601	FAD	C2A-N3A	3.78	1.38	1.32
2	A	601	FAD	C7M-C7	3.31	1.57	1.51
2	A	601	FAD	C4-N3	-3.26	1.32	1.38
4	A	603	TAM	C3-C6	-3.00	1.46	1.52
2	A	601	FAD	O4B-C1B	2.59	1.44	1.41
2	A	601	FAD	C5X-N5	-2.54	1.34	1.39
2	A	601	FAD	C6-C5X	-2.49	1.36	1.40
2	A	601	FAD	O4-C4	2.44	1.28	1.23
2	A	601	FAD	C9-C8	-2.30	1.36	1.39
2	A	601	FAD	PA-O1A	-2.18	1.43	1.50
2	A	601	FAD	C6-C7	2.16	1.42	1.39
3	A	602	SO4	O2-S	2.10	1.57	1.46

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	603	TAM	C3-C-N	-14.24	67.51	108.09
4	A	603	TAM	C3-C-C1	6.70	122.32	110.50
4	A	603	TAM	O6-C6-C3	-6.16	94.93	111.39
4	A	603	TAM	C2-C-C1	5.01	119.34	110.50
4	A	603	TAM	C1-C-N	-4.37	95.64	108.09
4	A	603	TAM	C3-C-C2	3.95	117.47	110.50
2	A	601	FAD	O4-C4-C4X	-3.90	116.25	126.60
2	A	601	FAD	C10-N1-C2	3.68	124.26	116.90
4	A	603	TAM	C2-C-N	-3.65	97.69	108.09
2	A	601	FAD	C1B-N9A-C4A	-3.50	120.49	126.64
2	A	601	FAD	N3A-C2A-N1A	-3.46	123.27	128.68
2	A	601	FAD	O4B-C1B-C2B	-3.35	102.03	106.93
2	A	601	FAD	O2'-C2'-C1'	-3.32	101.77	109.80
2	A	601	FAD	C4-C4X-N5	3.16	122.73	118.23
2	A	601	FAD	C4X-C4-N3	3.01	120.82	113.19
2	A	601	FAD	O2P-P-O1P	3.00	127.06	112.24
2	A	601	FAD	C6-C5X-N5	2.99	123.73	118.51
2	A	601	FAD	N6A-C6A-N1A	2.71	124.20	118.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	O3'-C3'-C4'	-2.55	102.65	108.81
2	A	601	FAD	C4'-C3'-C2'	2.48	118.52	113.36
2	A	601	FAD	C4-C4X-C10	-2.44	112.70	116.79
2	A	601	FAD	C9A-C5X-N5	-2.27	119.96	122.43
2	A	601	FAD	O2'-C2'-C3'	-2.19	103.77	109.10
2	A	601	FAD	C6-C5X-C9A	-2.16	115.89	118.94
2	A	601	FAD	C4X-C10-N1	-2.02	120.04	124.73

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	FAD	N10-C1'-C2'-O2'
2	A	601	FAD	N10-C1'-C2'-C3'
4	A	603	TAM	C2-C-C1-C4
4	A	603	TAM	N-C-C1-C4
4	A	603	TAM	C1-C-C2-C5
4	A	603	TAM	C3-C-C2-C5
4	A	603	TAM	C1-C-C3-C6
4	A	603	TAM	C-C3-C6-O6
2	A	601	FAD	O3'-C3'-C4'-C5'
2	A	601	FAD	C2'-C3'-C4'-C5'
4	A	603	TAM	C-C2-C5-O5
2	A	601	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	TAM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/549 (95%)	-0.14	6 (1%) 80 82	13, 26, 45, 84	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	ALA	7.3
1	A	177	GLU	4.5
1	A	241	ASP	3.0
1	A	245	GLU	2.4
1	A	182	LYS	2.3
1	A	240	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no monosaccharides in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TAM	A	603	11/11	0.90	0.14	35,46,52,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	602	5/5	0.98	0.14	39,40,41,42	0
2	FAD	A	601	53/53	0.99	0.10	13,16,23,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.