



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2024 – 02:57 PM EDT

PDB ID : 4JO7  
Title : Crystal structure of the human Nup49CCS2+3\* Nup57CCS3\* complex with 2:2 stoichiometry  
Authors : Stuwe, T.; Bley, C.J.; Mayo, D.J.; Hoelz, A.  
Deposited on : 2013-03-17  
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	40	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 17%, orange 82%, yellow 88%, green 92%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>17%</span> <span>82%</span> <span>88%</span> <span>92%</span> </div> </div>
1	D	40	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 8%, green 92%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>0%</span> <span>0%</span> <span>8%</span> <span>92%</span> </div> </div>
1	F	40	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 15%, green 85%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>0%</span> <span>0%</span> <span>15%</span> <span>85%</span> </div> </div>
1	H	40	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 0%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>0%</span> <span>0%</span> <span>0%</span> <span>98%</span> </div> </div>
2	A	89	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 17%, orange 82%, yellow 82%, green 98%, grey 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>17%</span> <span>82%</span> <span>82%</span> <span>98%</span> <span>11%</span> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	89	<div> <div>10%</div> <div>87%</div> <div>9%</div> <div>•</div> </div>
2	E	89	<div> <div>10%</div> <div>84%</div> <div>7%</div> <div>9%</div> </div>
2	G	89	<div> <div>11%</div> <div>74%</div> <div>17%</div> <div>•</div> <div>8%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8763 atoms, of which 4171 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoporin p54.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	40	Total	C	H	N	O	0	1	0
			684	214	348	55	67			
1	H	40	Total	C	H	N	O	0	0	0
			676	212	343	55	66			
1	D	40	Total	C	H	N	O	0	0	0
			676	212	343	55	66			
1	F	40	Total	C	H	N	O	0	0	0
			676	212	343	55	66			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	452	SER	-	EXPRESSION TAG	UNP Q7Z3B4
H	452	SER	-	EXPRESSION TAG	UNP Q7Z3B4
D	452	SER	-	EXPRESSION TAG	UNP Q7Z3B4
F	452	SER	-	EXPRESSION TAG	UNP Q7Z3B4

- Molecule 2 is a protein called Nucleoporin p58/p45.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	E	81	Total	C	H	N	O	Se	0	1	0
			1370	440	688	116	123	3			
2	G	82	Total	C	H	N	O	Se	0	2	0
			1391	449	693	117	129	3			
2	A	85	Total	C	H	N	O	Se	0	2	0
			1429	460	713	120	133	3			
2	C	85	Total	C	H	N	O	Se	0	0	0
			1407	450	700	122	132	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	328	SER	-	EXPRESSION TAG	UNP Q9BVL2
G	328	SER	-	EXPRESSION TAG	UNP Q9BVL2
A	328	SER	-	EXPRESSION TAG	UNP Q9BVL2
C	328	SER	-	EXPRESSION TAG	UNP Q9BVL2


- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	41	Total O 41 41	0	0
3	E	69	Total O 69 69	0	0
3	H	49	Total O 49 49	0	0
3	D	40	Total O 40 40	0	0
3	G	56	Total O 56 56	0	0
3	A	71	Total O 71 71	0	0
3	C	81	Total O 81 81	0	0
3	F	47	Total O 47 47	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoporin p54

Chain B:  88% 12%



- Molecule 1: Nucleoporin p54

Chain H:  98% .





- Molecule 1: Nucleoporin p54

Chain D:  92% 8%





- Molecule 1: Nucleoporin p54

Chain F:  85% 15%




- Molecule 2: Nucleoporin p58/p45



Chain E:  10% 84% 7% 9%





- Molecule 2: Nucleoporin p58/p45



• Molecule 2: Nucleoporin p58/p45



• Molecule 2: Nucleoporin p58/p45



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.69Å 47.66Å 71.32Å 88.87° 83.37° 81.42°	Depositor
Resolution (Å)	47.12 – 1.75 47.12 – 1.75	Depositor EDS
% Data completeness (in resolution range)	94.3 (47.12-1.75) 88.2 (47.12-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 1.75Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, $R_{free}$	0.186 , 0.214 0.189 , 0.216	Depositor DCC
$R_{free}$ test set	1965 reflections (3.35%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtriage
Anisotropy	0.426	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.46 , 55.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8763	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.83	0/342	0.52	0/457
1	D	0.39	0/336	0.42	0/449
1	F	0.40	0/336	0.41	0/449
1	H	0.55	0/336	0.37	0/449
2	A	0.38	0/732	0.46	0/981
2	C	0.32	0/716	0.39	0/960
2	E	0.35	0/694	0.39	0/927
2	G	0.35	0/711	0.40	0/952
All	All	0.43	0/4203	0.42	0/5624

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	336	348	346	5	0
1	D	333	343	341	3	0
1	F	333	343	341	7	0
1	H	333	343	341	0	0
2	A	716	713	710	13	0
2	C	707	700	697	10	0
2	E	682	688	685	9	0
2	G	698	693	689	11	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	71	0	0	2	0
3	B	41	0	0	2	0
3	C	81	0	0	2	0
3	D	40	0	0	2	0
3	E	69	0	0	1	0
3	F	47	0	0	1	0
3	G	56	0	0	0	0
3	H	49	0	0	1	0
All	All	4592	4171	4150	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:465:GLN:NE2	3:D:521:HOH:O	2.29	0.66
2:E:379[B]:LYS:NZ	3:E:554:HOH:O	2.29	0.64
1:F:484:ASP:O	1:F:488:ILE:HG12	1.98	0.64
2:A:378:GLN:NE2	3:A:569:HOH:O	2.34	0.60
2:G:366:SER:HA	2:G:367:HIS:C	2.23	0.58

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	39/40 (98%)	39 (100%)	0	0	100	100
1	D	38/40 (95%)	37 (97%)	1 (3%)	0	100	100
1	F	38/40 (95%)	37 (97%)	1 (3%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	38/40 (95%)	37 (97%)	1 (3%)	0	100	100
2	A	83/89 (93%)	82 (99%)	1 (1%)	0	100	100
2	C	83/89 (93%)	83 (100%)	0	0	100	100
2	E	78/89 (88%)	78 (100%)	0	0	100	100
2	G	80/89 (90%)	80 (100%)	0	0	100	100
All	All	477/516 (92%)	473 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	39/38 (103%)	38 (97%)	1 (3%)	46	23
1	D	38/38 (100%)	38 (100%)	0	100	100
1	F	38/38 (100%)	37 (97%)	1 (3%)	46	23
1	H	38/38 (100%)	37 (97%)	1 (3%)	46	23
2	A	78/75 (104%)	73 (94%)	5 (6%)	17	3
2	C	76/75 (101%)	75 (99%)	1 (1%)	69	54
2	E	73/75 (97%)	73 (100%)	0	100	100
2	G	76/75 (101%)	72 (95%)	4 (5%)	22	5
All	All	456/452 (101%)	443 (97%)	13 (3%)	42	19

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	358	HIS
2	A	359	LEU
1	F	486	GLU
2	A	371	GLN
2	C	367	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	E	392	GLN
2	A	367	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	40/40 (100%)	-0.05	0 100 100	10, 17, 28, 34	0
1	D	40/40 (100%)	-0.09	0 100 100	9, 16, 28, 36	0
1	F	40/40 (100%)	-0.03	0 100 100	9, 17, 29, 32	0
1	H	40/40 (100%)	0.02	0 100 100	9, 16, 32, 33	0
2	A	82/89 (92%)	0.88	15 (18%) 1 1	9, 20, 58, 76	0
2	C	82/89 (92%)	0.44	9 (10%) 5 7	8, 18, 47, 59	0
2	E	78/89 (87%)	0.84	9 (11%) 4 6	8, 17, 54, 68	0
2	G	79/89 (88%)	0.65	10 (12%) 3 5	8, 19, 61, 83	0
All	All	481/516 (93%)	0.45	43 (8%) 9 12	8, 17, 53, 83	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	360	ALA	28.8
2	A	362	GLN	7.1
2	E	368	ILE	6.4
2	G	367	HIS	6.4
2	A	361	THR	6.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.