



Full wwPDB NMR Structure Validation Report ⓘ

Jun 15, 2024 – 08:21 PM EDT

PDB ID : 2MYT
BMRB ID : 17077
Title : An arsenate reductase in the intermediate state
Authors : Jin, C.; Yu, C.; Hu, C.; Hu, Y.
Deposited on : 2015-01-30

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	134	

2 Ensemble composition and analysis

This entry contains 20 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:131 (131)	0.81	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	3, 8, 9, 10, 11, 12, 14, 15, 18, 19, 20
2	1, 4, 5, 6, 7, 16
3	13, 17
Single-model clusters	2

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1997 atoms, of which 990 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Glutaredoxin arsenate reductase.

Mol	Chain	Residues	Atoms						Trace
1	A	131	Total	C	H	N	O	S	0
			1997	626	990	171	203	7	

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P74313
A	-1	SER	-	EXPRESSION TAG	UNP P74313
A	0	HIS	-	EXPRESSION TAG	UNP P74313
A	13	SER	CYS	ENGINEERED MUTATION	UNP P74313
A	35	SER	CYS	ENGINEERED MUTATION	UNP P74313
A	82	SER	CYS	ENGINEERED MUTATION	UNP P74313

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Glutaredoxin arsenate reductase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Glutaredoxin arsenate reductase




4.2.2 Score per residue for model 2


- Molecule 1: Glutaredoxin arsenate reductase



4.2.3 Score per residue for model 3

- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  90% 8% .



4.2.4 Score per residue for model 4


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  88% 9% ..



4.2.5 Score per residue for model 5


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  85% 12% ..



4.2.6 Score per residue for model 6


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  87% 9% ..



4.2.7 Score per residue for model 7


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  84% 13% ..



4.2.8 Score per residue for model 8


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  86% 12% .



4.2.9 Score per residue for model 9


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  87% 10% ..



4.2.10 Score per residue for model 10 (medoid)


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  85% 11% ..



4.2.11 Score per residue for model 11


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  86% 12% .



4.2.12 Score per residue for model 12


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  86% 10% ..




4.2.13 Score per residue for model 13

- Molecule 1: Glutaredoxin arsenate reductase




4.2.14 Score per residue for model 14

- Molecule 1: Glutaredoxin arsenate reductase




4.2.15 Score per residue for model 15

- Molecule 1: Glutaredoxin arsenate reductase




4.2.16 Score per residue for model 16

- Molecule 1: Glutaredoxin arsenate reductase




4.2.17 Score per residue for model 17

- Molecule 1: Glutaredoxin arsenate reductase



4.2.18 Score per residue for model 18


- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  88% 10% .



4.2.19 Score per residue for model 19

- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  85% 12% ..



4.2.20 Score per residue for model 20

- Molecule 1: Glutaredoxin arsenate reductase

Chain A:  87% 10% ..



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AMBER	refinement	
CYANA	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1525
Number of shifts mapped to atoms	1524
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.70±0.01	0±0/1022 (0.0± 0.0%)	1.06±0.02	3±2/1378 (0.2± 0.1%)
All	All	0.70	0/20440 (0.0%)	1.06	58/27560 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.1±0.3
All	All	0	2

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	10	ARG	NE-CZ-NH1	8.31	124.46	120.30	10	11
1	A	122	ARG	NE-CZ-NH1	7.09	123.84	120.30	12	9
1	A	41	ARG	NE-CZ-NH1	6.91	123.75	120.30	10	5
1	A	14	ARG	NE-CZ-NH1	6.72	123.66	120.30	10	11
1	A	10	ARG	NE-CZ-NH2	-6.39	117.11	120.30	8	4
1	A	116	ARG	NE-CZ-NH1	6.24	123.42	120.30	18	13
1	A	14	ARG	NE-CZ-NH2	-6.23	117.19	120.30	20	2
1	A	113	ARG	NE-CZ-NH1	5.53	123.07	120.30	8	2
1	A	14	ARG	CD-NE-CZ	5.15	130.81	123.60	20	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	10	ARG	Sidechain	2

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1007	990	992	1±0
All	All	20140	19800	19840	20

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:75:VAL:HG22	1:A:95:ILE:HB	0.56	1.78	8	18
1:A:7:VAL:HG11	1:A:65:ILE:HD12	0.42	1.92	5	2

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/134 (96%)	120±2 (93±2%)	8±2 (6±2%)	1±1 (1±1%)	21	69
All	All	2580/2680 (96%)	2392 (93%)	166 (6%)	22 (1%)	21	69

All 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	ARG	4
1	A	41	ARG	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	13	SER	3
1	A	62	SER	2
1	A	84	VAL	1
1	A	103	ASP	1
1	A	105	ASP	1
1	A	11	ASN	1
1	A	83	GLY	1
1	A	60	GLN	1
1	A	81	GLY	1
1	A	17	MET	1
1	A	39	SER	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/115 (98%)	103±2 (91±1%)	10±2 (9±1%)	13	60
All	All	2260/2300 (98%)	2060 (91%)	200 (9%)	13	60

All 35 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	17	MET	20
1	A	49	MET	18
1	A	21	PHE	16
1	A	86	LEU	16
1	A	2	LYS	15
1	A	71	ASP	14
1	A	43	HIS	11
1	A	112	PHE	11
1	A	122	ARG	11
1	A	92	THR	10
1	A	67	ASN	10
1	A	50	MET	6
1	A	7	VAL	5
1	A	103	ASP	5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	72	ASP	3
1	A	8	CYS	3
1	A	85	ASN	3
1	A	102	GLU	2
1	A	19	GLU	2
1	A	23	LYS	2
1	A	63	ASP	2
1	A	101	LEU	2
1	A	14	ARG	1
1	A	55	ILE	1
1	A	24	THR	1
1	A	105	ASP	1
1	A	5	MET	1
1	A	16	GLN	1
1	A	37	LEU	1
1	A	61	THR	1
1	A	107	GLN	1
1	A	60	GLN	1
1	A	79	LEU	1
1	A	66	GLU	1
1	A	80	CYS	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 88% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1525
Number of shifts mapped to atoms	1524
Number of unparsed shifts	0
Number of shifts with mapping errors	1
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	17

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 1 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	1	MET	H1	8.31	.	.

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	130	-0.44 ± 0.13	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	120	0.12 ± 0.11	None needed (< 0.5 ppm)
$^{13}\text{C}'$	128	-0.15 ± 0.14	None needed (< 0.5 ppm)
^{15}N	124	-0.56 ± 0.29	None needed (imprecise)

7.1.3 Completeness of resonance assignments ⓘ

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 1524 atoms were assigned a chemical shift out of a possible 1728. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	645/655 (98%)	263/267 (99%)	258/262 (98%)	124/126 (98%)
Sidechain	834/983 (85%)	560/638 (88%)	263/309 (85%)	11/36 (31%)
Aromatic	45/90 (50%)	23/45 (51%)	20/42 (48%)	2/3 (67%)
Overall	1524/1728 (88%)	846/950 (89%)	541/613 (88%)	137/165 (83%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1524 atoms were assigned a chemical shift out of a possible 1728. 0 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹H	¹³C	¹⁵N
Backbone	645/655 (98%)	263/267 (99%)	258/262 (98%)	124/126 (98%)
Sidechain	834/983 (85%)	560/638 (88%)	263/309 (85%)	11/36 (31%)
Aromatic	45/90 (50%)	23/45 (51%)	20/42 (48%)	2/3 (67%)
Overall	1524/1728 (88%)	846/950 (89%)	541/613 (88%)	137/165 (83%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	82	SER	CB	27.58	56.28 – 71.32	-24.1
1	A	13	SER	CB	28.66	56.28 – 71.32	-23.4
1	A	35	SER	CB	31.44	56.28 – 71.32	-21.5
1	A	122	ARG	HB3	-0.57	0.43 – 3.11	-8.7
1	A	112	PHE	CE2	139.56	124.80 – 136.72	7.4
1	A	77	ILE	HG21	-1.09	-0.56 – 2.11	-7.0
1	A	77	ILE	HG22	-1.09	-0.56 – 2.11	-7.0
1	A	77	ILE	HG23	-1.09	-0.56 – 2.11	-7.0
1	A	70	ALA	HB1	-0.28	0.14 – 2.58	-6.7
1	A	70	ALA	HB2	-0.28	0.14 – 2.58	-6.7
1	A	70	ALA	HB3	-0.28	0.14 – 2.58	-6.7
1	A	68	PHE	CZ	120.38	121.82 – 136.66	-6.0
1	A	35	SER	HB3	2.24	2.49 – 5.20	-5.9
1	A	35	SER	HB2	2.43	2.61 – 5.13	-5.7

Continued on next page...

Continued from previous page...

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	45	THR	H	11.62	5.19 – 11.27	5.6
1	A	70	ALA	HA	1.98	2.13 – 6.34	-5.4
1	A	104	PRO	HB3	0.16	0.25 – 3.76	-5.3

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

