



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 04:54 AM EDT

PDB ID : 5ODN
Title : Salinibacter ruber Single-Strand Binding protein
Authors : Pierechod, M.; Rothweiler, U.
Deposited on : 2017-07-06
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	
1	B	196	
1	C	196	
1	D	196	
1	E	196	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	196	
1	G	196	
1	H	196	
2	I	4	
2	J	4	
2	K	4	
2	M	4	
2	N	4	
2	Q	4	
3	L	3	
3	P	3	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7084 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single-stranded DNA-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	0	0
			778	481	134	157	6			
1	B	106	Total	C	N	O	S	0	0	0
			826	511	139	170	6			
1	C	102	Total	C	N	O	S	0	0	0
			784	485	133	160	6			
1	D	106	Total	C	N	O	S	0	0	0
			831	517	142	166	6			
1	E	105	Total	C	N	O	S	0	0	0
			820	507	137	170	6			
1	F	105	Total	C	N	O	S	0	0	0
			808	502	132	168	6			
1	G	102	Total	C	N	O	S	0	0	0
			785	489	131	159	6			
1	H	99	Total	C	N	O	S	0	0	0
			778	481	134	157	6			

There are 224 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	initiating methionine	UNP Q2S565
A	-26	ASN	-	expression tag	UNP Q2S565
A	-25	HIS	-	expression tag	UNP Q2S565
A	-24	LYS	-	expression tag	UNP Q2S565
A	-23	VAL	-	expression tag	UNP Q2S565
A	-22	HIS	-	expression tag	UNP Q2S565
A	-21	HIS	-	expression tag	UNP Q2S565
A	-20	HIS	-	expression tag	UNP Q2S565
A	-19	HIS	-	expression tag	UNP Q2S565
A	-18	HIS	-	expression tag	UNP Q2S565
A	-17	HIS	-	expression tag	UNP Q2S565
A	-16	MET	-	expression tag	UNP Q2S565
A	-15	GLU	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	LEU	-	expression tag	UNP Q2S565
A	-13	GLY	-	expression tag	UNP Q2S565
A	-12	THR	-	expression tag	UNP Q2S565
A	-11	LEU	-	expression tag	UNP Q2S565
A	-10	GLU	-	expression tag	UNP Q2S565
A	-9	GLY	-	expression tag	UNP Q2S565
A	-8	SER	-	expression tag	UNP Q2S565
A	-7	GLU	-	expression tag	UNP Q2S565
A	-6	ASN	-	expression tag	UNP Q2S565
A	-5	LEU	-	expression tag	UNP Q2S565
A	-4	TYR	-	expression tag	UNP Q2S565
A	-3	PHE	-	expression tag	UNP Q2S565
A	-2	GLN	-	expression tag	UNP Q2S565
A	-1	GLY	-	expression tag	UNP Q2S565
A	0	ALA	-	expression tag	UNP Q2S565
B	-27	MET	-	initiating methionine	UNP Q2S565
B	-26	ASN	-	expression tag	UNP Q2S565
B	-25	HIS	-	expression tag	UNP Q2S565
B	-24	LYS	-	expression tag	UNP Q2S565
B	-23	VAL	-	expression tag	UNP Q2S565
B	-22	HIS	-	expression tag	UNP Q2S565
B	-21	HIS	-	expression tag	UNP Q2S565
B	-20	HIS	-	expression tag	UNP Q2S565
B	-19	HIS	-	expression tag	UNP Q2S565
B	-18	HIS	-	expression tag	UNP Q2S565
B	-17	HIS	-	expression tag	UNP Q2S565
B	-16	MET	-	expression tag	UNP Q2S565
B	-15	GLU	-	expression tag	UNP Q2S565
B	-14	LEU	-	expression tag	UNP Q2S565
B	-13	GLY	-	expression tag	UNP Q2S565
B	-12	THR	-	expression tag	UNP Q2S565
B	-11	LEU	-	expression tag	UNP Q2S565
B	-10	GLU	-	expression tag	UNP Q2S565
B	-9	GLY	-	expression tag	UNP Q2S565
B	-8	SER	-	expression tag	UNP Q2S565
B	-7	GLU	-	expression tag	UNP Q2S565
B	-6	ASN	-	expression tag	UNP Q2S565
B	-5	LEU	-	expression tag	UNP Q2S565
B	-4	TYR	-	expression tag	UNP Q2S565
B	-3	PHE	-	expression tag	UNP Q2S565
B	-2	GLN	-	expression tag	UNP Q2S565
B	-1	GLY	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	ALA	-	expression tag	UNP Q2S565
C	-27	MET	-	initiating methionine	UNP Q2S565
C	-26	ASN	-	expression tag	UNP Q2S565
C	-25	HIS	-	expression tag	UNP Q2S565
C	-24	LYS	-	expression tag	UNP Q2S565
C	-23	VAL	-	expression tag	UNP Q2S565
C	-22	HIS	-	expression tag	UNP Q2S565
C	-21	HIS	-	expression tag	UNP Q2S565
C	-20	HIS	-	expression tag	UNP Q2S565
C	-19	HIS	-	expression tag	UNP Q2S565
C	-18	HIS	-	expression tag	UNP Q2S565
C	-17	HIS	-	expression tag	UNP Q2S565
C	-16	MET	-	expression tag	UNP Q2S565
C	-15	GLU	-	expression tag	UNP Q2S565
C	-14	LEU	-	expression tag	UNP Q2S565
C	-13	GLY	-	expression tag	UNP Q2S565
C	-12	THR	-	expression tag	UNP Q2S565
C	-11	LEU	-	expression tag	UNP Q2S565
C	-10	GLU	-	expression tag	UNP Q2S565
C	-9	GLY	-	expression tag	UNP Q2S565
C	-8	SER	-	expression tag	UNP Q2S565
C	-7	GLU	-	expression tag	UNP Q2S565
C	-6	ASN	-	expression tag	UNP Q2S565
C	-5	LEU	-	expression tag	UNP Q2S565
C	-4	TYR	-	expression tag	UNP Q2S565
C	-3	PHE	-	expression tag	UNP Q2S565
C	-2	GLN	-	expression tag	UNP Q2S565
C	-1	GLY	-	expression tag	UNP Q2S565
C	0	ALA	-	expression tag	UNP Q2S565
D	-27	MET	-	initiating methionine	UNP Q2S565
D	-26	ASN	-	expression tag	UNP Q2S565
D	-25	HIS	-	expression tag	UNP Q2S565
D	-24	LYS	-	expression tag	UNP Q2S565
D	-23	VAL	-	expression tag	UNP Q2S565
D	-22	HIS	-	expression tag	UNP Q2S565
D	-21	HIS	-	expression tag	UNP Q2S565
D	-20	HIS	-	expression tag	UNP Q2S565
D	-19	HIS	-	expression tag	UNP Q2S565
D	-18	HIS	-	expression tag	UNP Q2S565
D	-17	HIS	-	expression tag	UNP Q2S565
D	-16	MET	-	expression tag	UNP Q2S565
D	-15	GLU	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	LEU	-	expression tag	UNP Q2S565
D	-13	GLY	-	expression tag	UNP Q2S565
D	-12	THR	-	expression tag	UNP Q2S565
D	-11	LEU	-	expression tag	UNP Q2S565
D	-10	GLU	-	expression tag	UNP Q2S565
D	-9	GLY	-	expression tag	UNP Q2S565
D	-8	SER	-	expression tag	UNP Q2S565
D	-7	GLU	-	expression tag	UNP Q2S565
D	-6	ASN	-	expression tag	UNP Q2S565
D	-5	LEU	-	expression tag	UNP Q2S565
D	-4	TYR	-	expression tag	UNP Q2S565
D	-3	PHE	-	expression tag	UNP Q2S565
D	-2	GLN	-	expression tag	UNP Q2S565
D	-1	GLY	-	expression tag	UNP Q2S565
D	0	ALA	-	expression tag	UNP Q2S565
E	-27	MET	-	initiating methionine	UNP Q2S565
E	-26	ASN	-	expression tag	UNP Q2S565
E	-25	HIS	-	expression tag	UNP Q2S565
E	-24	LYS	-	expression tag	UNP Q2S565
E	-23	VAL	-	expression tag	UNP Q2S565
E	-22	HIS	-	expression tag	UNP Q2S565
E	-21	HIS	-	expression tag	UNP Q2S565
E	-20	HIS	-	expression tag	UNP Q2S565
E	-19	HIS	-	expression tag	UNP Q2S565
E	-18	HIS	-	expression tag	UNP Q2S565
E	-17	HIS	-	expression tag	UNP Q2S565
E	-16	MET	-	expression tag	UNP Q2S565
E	-15	GLU	-	expression tag	UNP Q2S565
E	-14	LEU	-	expression tag	UNP Q2S565
E	-13	GLY	-	expression tag	UNP Q2S565
E	-12	THR	-	expression tag	UNP Q2S565
E	-11	LEU	-	expression tag	UNP Q2S565
E	-10	GLU	-	expression tag	UNP Q2S565
E	-9	GLY	-	expression tag	UNP Q2S565
E	-8	SER	-	expression tag	UNP Q2S565
E	-7	GLU	-	expression tag	UNP Q2S565
E	-6	ASN	-	expression tag	UNP Q2S565
E	-5	LEU	-	expression tag	UNP Q2S565
E	-4	TYR	-	expression tag	UNP Q2S565
E	-3	PHE	-	expression tag	UNP Q2S565
E	-2	GLN	-	expression tag	UNP Q2S565
E	-1	GLY	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	0	ALA	-	expression tag	UNP Q2S565
F	-27	MET	-	initiating methionine	UNP Q2S565
F	-26	ASN	-	expression tag	UNP Q2S565
F	-25	HIS	-	expression tag	UNP Q2S565
F	-24	LYS	-	expression tag	UNP Q2S565
F	-23	VAL	-	expression tag	UNP Q2S565
F	-22	HIS	-	expression tag	UNP Q2S565
F	-21	HIS	-	expression tag	UNP Q2S565
F	-20	HIS	-	expression tag	UNP Q2S565
F	-19	HIS	-	expression tag	UNP Q2S565
F	-18	HIS	-	expression tag	UNP Q2S565
F	-17	HIS	-	expression tag	UNP Q2S565
F	-16	MET	-	expression tag	UNP Q2S565
F	-15	GLU	-	expression tag	UNP Q2S565
F	-14	LEU	-	expression tag	UNP Q2S565
F	-13	GLY	-	expression tag	UNP Q2S565
F	-12	THR	-	expression tag	UNP Q2S565
F	-11	LEU	-	expression tag	UNP Q2S565
F	-10	GLU	-	expression tag	UNP Q2S565
F	-9	GLY	-	expression tag	UNP Q2S565
F	-8	SER	-	expression tag	UNP Q2S565
F	-7	GLU	-	expression tag	UNP Q2S565
F	-6	ASN	-	expression tag	UNP Q2S565
F	-5	LEU	-	expression tag	UNP Q2S565
F	-4	TYR	-	expression tag	UNP Q2S565
F	-3	PHE	-	expression tag	UNP Q2S565
F	-2	GLN	-	expression tag	UNP Q2S565
F	-1	GLY	-	expression tag	UNP Q2S565
F	0	ALA	-	expression tag	UNP Q2S565
G	-27	MET	-	initiating methionine	UNP Q2S565
G	-26	ASN	-	expression tag	UNP Q2S565
G	-25	HIS	-	expression tag	UNP Q2S565
G	-24	LYS	-	expression tag	UNP Q2S565
G	-23	VAL	-	expression tag	UNP Q2S565
G	-22	HIS	-	expression tag	UNP Q2S565
G	-21	HIS	-	expression tag	UNP Q2S565
G	-20	HIS	-	expression tag	UNP Q2S565
G	-19	HIS	-	expression tag	UNP Q2S565
G	-18	HIS	-	expression tag	UNP Q2S565
G	-17	HIS	-	expression tag	UNP Q2S565
G	-16	MET	-	expression tag	UNP Q2S565
G	-15	GLU	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-14	LEU	-	expression tag	UNP Q2S565
G	-13	GLY	-	expression tag	UNP Q2S565
G	-12	THR	-	expression tag	UNP Q2S565
G	-11	LEU	-	expression tag	UNP Q2S565
G	-10	GLU	-	expression tag	UNP Q2S565
G	-9	GLY	-	expression tag	UNP Q2S565
G	-8	SER	-	expression tag	UNP Q2S565
G	-7	GLU	-	expression tag	UNP Q2S565
G	-6	ASN	-	expression tag	UNP Q2S565
G	-5	LEU	-	expression tag	UNP Q2S565
G	-4	TYR	-	expression tag	UNP Q2S565
G	-3	PHE	-	expression tag	UNP Q2S565
G	-2	GLN	-	expression tag	UNP Q2S565
G	-1	GLY	-	expression tag	UNP Q2S565
G	0	ALA	-	expression tag	UNP Q2S565
H	-27	MET	-	initiating methionine	UNP Q2S565
H	-26	ASN	-	expression tag	UNP Q2S565
H	-25	HIS	-	expression tag	UNP Q2S565
H	-24	LYS	-	expression tag	UNP Q2S565
H	-23	VAL	-	expression tag	UNP Q2S565
H	-22	HIS	-	expression tag	UNP Q2S565
H	-21	HIS	-	expression tag	UNP Q2S565
H	-20	HIS	-	expression tag	UNP Q2S565
H	-19	HIS	-	expression tag	UNP Q2S565
H	-18	HIS	-	expression tag	UNP Q2S565
H	-17	HIS	-	expression tag	UNP Q2S565
H	-16	MET	-	expression tag	UNP Q2S565
H	-15	GLU	-	expression tag	UNP Q2S565
H	-14	LEU	-	expression tag	UNP Q2S565
H	-13	GLY	-	expression tag	UNP Q2S565
H	-12	THR	-	expression tag	UNP Q2S565
H	-11	LEU	-	expression tag	UNP Q2S565
H	-10	GLU	-	expression tag	UNP Q2S565
H	-9	GLY	-	expression tag	UNP Q2S565
H	-8	SER	-	expression tag	UNP Q2S565
H	-7	GLU	-	expression tag	UNP Q2S565
H	-6	ASN	-	expression tag	UNP Q2S565
H	-5	LEU	-	expression tag	UNP Q2S565
H	-4	TYR	-	expression tag	UNP Q2S565
H	-3	PHE	-	expression tag	UNP Q2S565
H	-2	GLN	-	expression tag	UNP Q2S565
H	-1	GLY	-	expression tag	UNP Q2S565

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	0	ALA	-	expression tag	UNP Q2S565

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	4	Total	C	N	O	P	0	0	0
			80	40	8	28	4			
2	M	4	Total	C	N	O	P	0	0	0
			76	40	8	25	3			
2	J	4	Total	C	N	O	P	0	0	0
			80	40	8	28	4			
2	K	4	Total	C	N	O	P	0	0	0
			77	40	8	26	3			
2	N	4	Total	C	N	O	P	0	0	0
			64	30	6	24	4			
2	Q	4	Total	C	N	O	P	0	0	0
			77	40	8	26	3			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	3	Total	C	N	O	P	0	0	0
			57	30	6	19	2			
3	L	3	Total	C	N	O	P	0	0	0
			60	30	6	21	3			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	22	Total	O	0	0
			22	22		
4	B	10	Total	O	0	0
			10	10		
4	C	6	Total	O	0	0
			6	6		
4	D	14	Total	O	0	0
			14	14		
4	E	14	Total	O	0	0
			14	14		
4	F	9	Total	O	0	0
			9	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	10	Total	O	0	0
			10	10		
4	H	17	Total	O	0	0
			17	17		
4	K	1	Total	O	0	0
			1	1		



- Molecule 1: Single-stranded DNA-binding protein





- Molecule 1: Single-stranded DNA-binding protein




- Molecule 1: Single-stranded DNA-binding protein




- Molecule 1: Single-stranded DNA-binding protein




- Molecule 1: Single-stranded DNA-binding protein



- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')



- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')



- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')



- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')




There are no outlier residues recorded for this chain.

- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')



There are no outlier residues recorded for this chain.

- Molecule 2: DNA (5'-D(P*TP*TP*TP*T)-3')



- Molecule 3: DNA (5'-D(*TP*TP*T)-3')

Chain P:  33% 67%

T1
T2
T3

- Molecule 3: DNA (5'-D(*TP*TP*T)-3')

Chain L:  100% 67% 33%

T1
T2
T3

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	106.47Å 123.50Å 205.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.26 – 2.60 47.26 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.2 (47.26-2.60) 99.8 (47.26-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.190 , 0.243 0.189 , 0.242	Depositor DCC
R_{free} test set	2119 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7084	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/790	0.58	0/1067
1	B	0.42	0/839	0.55	0/1135
1	C	0.42	0/796	0.58	0/1078
1	D	0.41	0/845	0.57	0/1142
1	E	0.37	0/833	0.54	0/1127
1	F	0.37	0/821	0.56	0/1114
1	G	0.41	0/798	0.59	0/1081
1	H	0.42	0/790	0.57	0/1067
2	I	0.78	0/87	1.24	0/132
2	J	0.85	0/87	1.27	0/132
2	K	0.95	0/84	1.33	0/128
2	M	0.88	0/83	1.27	0/127
2	N	0.79	0/69	1.26	0/105
2	Q	0.73	0/84	1.25	0/128
3	L	0.65	0/65	1.29	0/98
3	P	0.78	0/62	1.32	0/94
All	All	0.46	0/7133	0.67	0/9755

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	778	0	736	14	0
1	B	826	0	769	8	0
1	C	784	0	728	12	0
1	D	831	0	778	17	0
1	E	820	0	763	10	0
1	F	808	0	739	21	0
1	G	785	0	734	16	0
1	H	778	0	736	13	0
2	I	80	0	49	1	0
2	J	80	0	49	0	0
2	K	77	0	50	0	0
2	M	76	0	47	1	0
2	N	64	0	36	0	0
2	Q	77	0	50	1	0
3	L	60	0	37	1	0
3	P	57	0	38	3	0
4	A	22	0	0	2	0
4	B	10	0	0	0	0
4	C	6	0	0	0	0
4	D	14	0	0	0	0
4	E	14	0	0	1	0
4	F	9	0	0	0	0
4	G	10	0	0	0	0
4	H	17	0	0	0	0
4	K	1	0	0	0	0
All	All	7084	0	6339	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:37:ASN:HB3	1:F:49:GLN:HE22	1.16	1.10
1:H:37:ASN:HD21	3:L:2:DT:H3	1.12	0.96
1:C:61:ARG:O	1:C:65:ILE:HD13	1.67	0.94
3:P:1:DT:H2''	3:P:2:DT:H5''	1.59	0.85
1:C:61:ARG:O	1:C:65:ILE:CD1	2.29	0.80

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/196 (48%)	95 (100%)	0	0	100	100
1	B	102/196 (52%)	101 (99%)	1 (1%)	0	100	100
1	C	98/196 (50%)	98 (100%)	0	0	100	100
1	D	102/196 (52%)	102 (100%)	0	0	100	100
1	E	101/196 (52%)	100 (99%)	1 (1%)	0	100	100
1	F	101/196 (52%)	99 (98%)	2 (2%)	0	100	100
1	G	98/196 (50%)	96 (98%)	2 (2%)	0	100	100
1	H	95/196 (48%)	95 (100%)	0	0	100	100
All	All	792/1568 (50%)	786 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/165 (50%)	83 (100%)	0	100	100
1	B	88/165 (53%)	88 (100%)	0	100	100
1	C	82/165 (50%)	82 (100%)	0	100	100
1	D	87/165 (53%)	84 (97%)	3 (3%)	37	63
1	E	88/165 (53%)	88 (100%)	0	100	100
1	F	85/165 (52%)	85 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	83/165 (50%)	82 (99%)	1 (1%)	71	87
1	H	83/165 (50%)	82 (99%)	1 (1%)	71	87
All	All	679/1320 (51%)	674 (99%)	5 (1%)	84	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	6	ASN
1	D	7	LYS
1	D	50	ASN
1	G	25	SER
1	H	3	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	6	ASN
1	F	49	GLN
1	H	75	GLN
1	F	103	GLN
1	H	37	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/196 (50%)	0.09	0 100 100	27, 43, 79, 101	0
1	B	106/196 (54%)	0.49	8 (7%) 14 10	32, 50, 96, 121	0
1	C	102/196 (52%)	0.31	3 (2%) 51 45	33, 51, 80, 99	0
1	D	106/196 (54%)	0.34	6 (5%) 23 18	35, 51, 84, 101	0
1	E	105/196 (53%)	0.46	6 (5%) 23 18	32, 58, 84, 103	0
1	F	105/196 (53%)	0.69	9 (8%) 10 7	35, 58, 89, 131	0
1	G	102/196 (52%)	0.67	13 (12%) 3 2	35, 57, 90, 108	0
1	H	99/196 (50%)	0.37	2 (2%) 65 60	31, 53, 76, 93	0
2	I	4/4 (100%)	0.61	1 (25%) 0 0	79, 85, 93, 109	0
2	J	4/4 (100%)	1.30	1 (25%) 0 0	84, 85, 98, 119	0
2	K	4/4 (100%)	0.54	0 100 100	64, 67, 82, 115	0
2	M	4/4 (100%)	-0.14	0 100 100	53, 55, 61, 98	0
2	N	4/4 (100%)	1.57	0 100 100	99, 100, 101, 118	0
2	Q	4/4 (100%)	0.67	1 (25%) 0 0	76, 80, 88, 104	0
3	L	3/3 (100%)	2.86	3 (100%) 0 0	112, 112, 116, 125	0
3	P	3/3 (100%)	1.12	0 100 100	87, 87, 108, 123	0
All	All	854/1598 (53%)	0.45	53 (6%) 20 15	27, 54, 91, 131	0

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	69	TYR	5.4
1	B	94	THR	5.1
1	D	-3	PHE	4.8
1	G	69	TYR	4.6
1	B	86	SER	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.