



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 08:07 AM EDT

PDB ID : 3P8K
Title : Crystal Structure of a putative carbon-nitrogen family hydrolase from *Staphylococcus aureus*
Authors : Gordon, R.D.; Qiu, W.; Battaile, K.; Lam, K.; Soloveychik, M.; Benetteraj, D.; Romanov, V.; Pai, E.F.; Chirgadze, N.Y.
Deposited on : 2010-10-14
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	
1	B	281	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	270	-	-	X	-
3	CL	A	271	-	-	X	-
4	PEG	A	264	-	X	X	-
4	PEG	A	265	-	-	X	-
4	PEG	B	262	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrolase, carbon-nitrogen family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	Se	0	3	0
			2173	1385	368	413	1	6			
1	B	268	Total	C	N	O	S	Se	0	3	0
			2175	1386	369	413	1	6			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	EXPRESSION TAG	UNP Q5HEG7
A	-18	GLY	-	EXPRESSION TAG	UNP Q5HEG7
A	-17	SER	-	EXPRESSION TAG	UNP Q5HEG7
A	-16	SER	-	EXPRESSION TAG	UNP Q5HEG7
A	-15	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-14	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-13	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-12	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-11	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-10	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	-9	SER	-	EXPRESSION TAG	UNP Q5HEG7
A	-8	SER	-	EXPRESSION TAG	UNP Q5HEG7
A	-7	GLY	-	EXPRESSION TAG	UNP Q5HEG7
A	-6	LEU	-	EXPRESSION TAG	UNP Q5HEG7
A	-5	VAL	-	EXPRESSION TAG	UNP Q5HEG7
A	-4	PRO	-	EXPRESSION TAG	UNP Q5HEG7
A	-3	ARG	-	EXPRESSION TAG	UNP Q5HEG7
A	-2	GLY	-	EXPRESSION TAG	UNP Q5HEG7
A	-1	SER	-	EXPRESSION TAG	UNP Q5HEG7
A	0	HIS	-	EXPRESSION TAG	UNP Q5HEG7
A	1	MSE	-	EXPRESSION TAG	UNP Q5HEG7
B	-19	MSE	-	EXPRESSION TAG	UNP Q5HEG7
B	-18	GLY	-	EXPRESSION TAG	UNP Q5HEG7
B	-17	SER	-	EXPRESSION TAG	UNP Q5HEG7
B	-16	SER	-	EXPRESSION TAG	UNP Q5HEG7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-14	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-13	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-12	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-11	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-10	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	-9	SER	-	EXPRESSION TAG	UNP Q5HEG7
B	-8	SER	-	EXPRESSION TAG	UNP Q5HEG7
B	-7	GLY	-	EXPRESSION TAG	UNP Q5HEG7
B	-6	LEU	-	EXPRESSION TAG	UNP Q5HEG7
B	-5	VAL	-	EXPRESSION TAG	UNP Q5HEG7
B	-4	PRO	-	EXPRESSION TAG	UNP Q5HEG7
B	-3	ARG	-	EXPRESSION TAG	UNP Q5HEG7
B	-2	GLY	-	EXPRESSION TAG	UNP Q5HEG7
B	-1	SER	-	EXPRESSION TAG	UNP Q5HEG7
B	0	HIS	-	EXPRESSION TAG	UNP Q5HEG7
B	1	MSE	-	EXPRESSION TAG	UNP Q5HEG7

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Cl	0	0
			4	4		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	286	Total 286	O 286	0	0
7	B	269	Total 269	O 269	0	0

i

- Molecule 1: Hydrolase, carbon-nitrogen family



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.58Å 61.98Å 155.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.96 – 1.70 19.68 – 1.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.96-1.70) 96.8 (19.68-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.70Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.197 , 0.230 0.201 , 0.235	Depositor DCC
R_{free} test set	3060 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4962	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PGE, TRS, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2225	0.67	0/3008
1	B	0.49	0/2227	0.68	0/3011
All	All	0.49	0/4452	0.68	0/6019

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2173	0	2127	21	0
1	B	2175	0	2128	22	0
2	A	8	0	12	1	0
3	A	4	0	0	5	0
4	A	14	0	17	13	0
4	B	7	0	10	6	0
5	A	20	0	28	0	0
6	A	6	0	8	1	0
7	A	286	0	0	1	0
7	B	269	0	0	0	0
All	All	4962	0	4330	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:250:PRO:HA	4:B:262:PEG:H32	1.64	0.78
4:A:265:PEG:H32	3:A:270:CL:CL	2.23	0.76
1:A:31:ASN:HD22	1:A:33:GLU:H	1.39	0.70
1:A:250:PRO:HA	4:A:264:PEG:H31	1.75	0.68
4:A:265:PEG:C3	3:A:270:CL:CL	2.79	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/281 (96%)	261 (97%)	7 (3%)	1 (0%)	34	18
1	B	269/281 (96%)	255 (95%)	12 (4%)	2 (1%)	22	8
All	All	538/562 (96%)	516 (96%)	19 (4%)	3 (1%)	25	11

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	CYS
1	B	146	CYS
1	B	118	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/242 (100%)	234 (97%)	7 (3%)	42	23
1	B	241/242 (100%)	239 (99%)	2 (1%)	81	74
All	All	482/484 (100%)	473 (98%)	9 (2%)	60	41

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	31	ASN
1	B	226	LEU
1	A	178[A]	GLN
1	A	178[B]	GLN
1	A	227	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	ASN
1	B	24	GLN
1	B	248	ASN
1	B	31	ASN
1	A	45	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	264	-	6,6,6	1.69	2 (33%)	5,5,5	1.18	1 (20%)
2	TRS	A	262	-	7,7,7	0.97	0	9,9,9	1.35	2 (22%)
4	PEG	B	262	-	6,6,6	1.68	2 (33%)	5,5,5	1.82	2 (40%)
4	PEG	A	265	-	6,6,6	1.28	1 (16%)	5,5,5	0.56	0
5	PGE	A	266	-	9,9,9	0.38	0	8,8,8	0.86	0
6	GOL	A	268	-	5,5,5	0.40	0	5,5,5	0.73	0
5	PGE	A	267	-	9,9,9	0.47	0	8,8,8	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	264	-	-	3/4/4/4	-
2	TRS	A	262	-	-	5/9/9/9	-
4	PEG	B	262	-	-	0/4/4/4	-
4	PEG	A	265	-	-	3/4/4/4	-
5	PGE	A	266	-	-	3/7/7/7	-
6	GOL	A	268	-	-	2/4/4/4	-
5	PGE	A	267	-	-	4/7/7/7	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	264	PEG	O2-C3	-2.76	1.30	1.42
4	A	265	PEG	C2-C1	-2.46	1.36	1.49
4	B	262	PEG	O1-C1	2.21	1.53	1.42
4	B	262	PEG	O2-C2	2.14	1.51	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	264	PEG	O2-C2	-2.04	1.33	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	262	TRS	O1-C1-C	2.81	118.70	110.88
4	B	262	PEG	O2-C2-C1	2.57	121.44	110.11
4	B	262	PEG	O4-C4-C3	2.41	125.98	111.82
2	A	262	TRS	C3-C-N	-2.25	102.42	108.17
4	A	264	PEG	O1-C1-C2	2.19	124.74	111.82

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	262	TRS	C2-C-C1-O1
2	A	262	TRS	N-C-C1-O1
6	A	268	GOL	O1-C1-C2-C3
5	A	267	PGE	O2-C3-C4-O3
4	A	264	PEG	O2-C3-C4-O4

There are no ring outliers.

5 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	264	PEG	5	0
2	A	262	TRS	1	0
4	B	262	PEG	6	0
4	A	265	PEG	8	0
6	A	268	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/281 (93%)	-0.12	6 (2%) 60 65	10, 18, 38, 61	0
1	B	262/281 (93%)	0.04	7 (2%) 54 58	10, 21, 40, 59	0
All	All	524/562 (93%)	-0.04	13 (2%) 57 61	10, 20, 40, 61	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	117	LEU	4.5
1	B	205	GLY	2.9
1	A	120	HIS	2.9
1	A	118	ARG	2.8
1	A	115	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PGE	A	267	10/10	0.62	0.37	53,64,70,72	0
5	PGE	A	266	10/10	0.63	0.27	47,59,65,71	0
4	PEG	A	265	7/7	0.69	0.25	35,37,39,42	0
2	TRS	A	262	8/8	0.73	0.22	37,38,42,42	0
6	GOL	A	268	6/6	0.81	0.15	43,44,44,45	0
4	PEG	B	262	7/7	0.85	0.24	22,22,26,31	0
4	PEG	A	264	7/7	0.90	0.25	16,18,24,30	0
3	CL	A	269	1/1	0.91	0.07	44,44,44,44	0
3	CL	A	270	1/1	0.94	0.08	36,36,36,36	0
3	CL	A	271	1/1	0.96	0.32	16,16,16,16	0
3	CL	A	263	1/1	0.99	0.03	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.