



# Full wwPDB X-ray Structure Validation Report i

May 23, 2020 – 08:38 am BST

PDB ID : 2R70  
Title : Crystal structure of infectious bursal disease virus VP1 polymerase, cocrystallized with an oligopeptide mimicking the VP3 C-terminus.  
Authors : Garriga, D.; Navarro, A.; Querol-Audi, J.; Abaitua, F.; Rodriguez, J.F.; Verdaguier, N.  
Deposited on : 2007-09-07  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	852	4%	73%	13%	.	12%

## 2 Entry composition [\(i\)](#)

There are 2 unique types of molecules in this entry. The entry contains 5990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INFECTIOUS BURSAL VIRUS VP1 POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	752	Total	C 5853	N 3747	O 994	S 1090	22	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	137	Total O 137 137	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INFECTIOUS BURSAL VIRUS VP1 POLYMERASE



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.80Å 123.80Å 361.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.70 34.27 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.9 (40.00-2.70) 94.0 (34.27-2.70)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	4.08 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
$R$ , $R_{free}$	0.221 , 0.256 0.241 , 0.271	Depositor DCC
$R_{free}$ test set	2195 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.7	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.7	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5990	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/5988	0.60	0/8136

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5853	0	5844	83	0
2	A	137	0	0	8	0
All	All	5990	0	5844	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:VAL:HG12	1:A:93:LEU:H	1.08	1.08
1:A:761:LYS:HB3	1:A:762:PRO:HA	1.42	1.02
1:A:92:VAL:HG12	1:A:93:LEU:N	1.72	1.01
1:A:92:VAL:CG1	1:A:93:LEU:H	1.78	0.96

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:SER:HB2	1:A:279:ASN:HD21	1.33	0.94
1:A:85:MET:O	1:A:92:VAL:HG22	1.69	0.91
1:A:301:LYS:HG2	1:A:301:LYS:O	1.70	0.90
1:A:459:ASN:O	1:A:797:THR:HG21	1.81	0.79
1:A:356:MET:CE	1:A:356:MET:HA	2.17	0.74
1:A:401:ALA:H	1:A:494:HIS:HD2	1.37	0.73
1:A:319:PHE:HD2	2:A:979:HOH:O	1.72	0.71
1:A:704:VAL:HG21	1:A:708:VAL:HG22	1.74	0.69
1:A:482:GLN:HG2	1:A:486:ASN:HD21	1.56	0.69
1:A:296:LYS:NZ	2:A:937:HOH:O	2.25	0.68
1:A:761:LYS:HB3	1:A:762:PRO:CA	2.22	0.67
1:A:589:ASP:HB3	1:A:592:ARG:HD2	1.77	0.66
1:A:600:PRO:HB3	1:A:622:ARG:HD2	1.77	0.65
1:A:319:PHE:CD2	2:A:979:HOH:O	2.48	0.65
1:A:92:VAL:CG1	1:A:93:LEU:N	2.43	0.64
1:A:294:GLY:HA2	1:A:299:ASN:ND2	2.16	0.61
1:A:65:ARG:HH11	1:A:476:GLN:HE21	1.47	0.61
1:A:793:ALA:O	1:A:797:THR:HG22	2.00	0.61
1:A:299:ASN:O	1:A:745:GLN:OE1	2.20	0.60
1:A:224:PRO:HB3	1:A:448:MET:HG3	1.84	0.59
1:A:356:MET:HA	1:A:356:MET:HE2	1.83	0.59
1:A:356:MET:HE3	1:A:356:MET:HA	1.85	0.59
1:A:347:MET:HA	1:A:350:MET:HE3	1.86	0.57
1:A:294:GLY:CA	1:A:299:ASN:ND2	2.67	0.57
1:A:482:GLN:CG	1:A:486:ASN:HD21	2.17	0.57
1:A:321:LYS:HA	2:A:979:HOH:O	2.04	0.57
1:A:425:THR:H	1:A:428:HIS:HD2	1.52	0.57
1:A:421:GLU:HA	2:A:883:HOH:O	2.05	0.56
1:A:482:GLN:HG2	1:A:486:ASN:ND2	2.20	0.56
1:A:482:GLN:NE2	2:A:883:HOH:O	2.40	0.55
1:A:170:MET:O	1:A:174:ASN:HB2	2.07	0.54
1:A:75:THR:OG1	1:A:95:PRO:O	2.23	0.54
1:A:318:LEU:HD22	1:A:336:ASN:HB3	1.89	0.53
1:A:401:ALA:H	1:A:494:HIS:CD2	2.24	0.53
1:A:299:ASN:O	1:A:745:GLN:NE2	2.41	0.53
1:A:93:LEU:O	1:A:94:LYS:O	2.27	0.53
1:A:339:SER:HB3	2:A:904:HOH:O	2.09	0.52
1:A:229:PRO:HG3	1:A:350:MET:CE	2.38	0.52
1:A:451:GLN:HA	1:A:451:GLN:NE2	2.25	0.52
1:A:761:LYS:HA	1:A:761:LYS:HE3	1.92	0.51
1:A:75:THR:HG1	1:A:95:PRO:C	2.14	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:LYS:HD3	1:A:337:ILE:HD11	1.92	0.51
1:A:450:ASN:HB3	1:A:452:THR:H	1.76	0.51
1:A:427:GLN:HA	1:A:430:GLN:HG2	1.93	0.50
1:A:230:SER:HB2	1:A:279:ASN:ND2	2.14	0.49
1:A:462:PRO:HB2	1:A:794:LEU:HD21	1.94	0.49
1:A:757:LEU:O	1:A:760:SER:HB3	2.13	0.49
1:A:180:LYS:HG3	1:A:480:TYR:CZ	2.48	0.49
1:A:65:ARG:HH11	1:A:476:GLN:NE2	2.11	0.48
1:A:595:CYS:SG	2:A:909:HOH:O	2.20	0.48
1:A:364:LEU:HD21	1:A:385:ILE:HD12	1.96	0.47
1:A:300:LYS:HB2	1:A:300:LYS:HE3	1.69	0.47
1:A:65:ARG:HD3	1:A:476:GLN:HE22	1.79	0.47
1:A:776:LEU:O	1:A:780:LEU:HG	2.15	0.46
1:A:716:VAL:O	1:A:720:LEU:HB2	2.16	0.45
1:A:65:ARG:H	1:A:476:GLN:HE22	1.65	0.45
1:A:75:THR:OG1	1:A:95:PRO:C	2.55	0.45
1:A:133:GLN:NE2	1:A:643:ASN:HD22	2.15	0.45
1:A:229:PRO:HG2	1:A:278:SER:HB2	1.99	0.44
1:A:301:LYS:HB3	1:A:301:LYS:HE2	1.24	0.44
1:A:616:GLN:HB2	1:A:652:LEU:HD21	2.00	0.44
1:A:425:THR:H	1:A:428:HIS:CD2	2.35	0.43
1:A:482:GLN:OE1	1:A:486:ASN:ND2	2.50	0.43
1:A:133:GLN:HE21	1:A:643:ASN:HD22	1.67	0.43
1:A:300:LYS:O	1:A:304:LEU:HG	2.18	0.43
1:A:37:TRP:CG	1:A:327:LYS:HE2	2.53	0.43
1:A:138:PHE:HB2	1:A:636:LEU:HD11	2.00	0.43
1:A:437:LEU:HD23	1:A:441:TRP:CD1	2.53	0.42
1:A:52:LYS:O	1:A:56:GLU:HB2	2.18	0.42
1:A:301:LYS:O	1:A:301:LYS:CG	2.51	0.42
1:A:295:THR:HA	1:A:296:LYS:HA	1.77	0.42
1:A:164:TYR:HA	1:A:323:GLU:O	2.19	0.41
1:A:424:CYS:HB3	1:A:482:GLN:HB2	2.03	0.41
1:A:34:PRO:HG2	1:A:159:ILE:HG22	2.02	0.41
1:A:199:PHE:CE1	1:A:430:GLN:HG3	2.56	0.40
1:A:170:MET:HG3	1:A:331:LEU:O	2.22	0.40
1:A:346:LEU:O	1:A:350:MET:HG3	2.21	0.40
1:A:482:GLN:CD	1:A:486:ASN:HD21	2.24	0.40
1:A:377:PRO:HG3	1:A:573:TRP:CE2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	740/852 (87%)	708 (96%)	25 (3%)	7 (1%)	17 40

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	419	LYS
1	A	767	ALA
1	A	468	SER
1	A	803	ALA
1	A	94	LYS
1	A	295	THR
1	A	685	VAL

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	628/730 (86%)	585 (93%)	43 (7%)	16 36

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	ARG
1	A	145	ASN
1	A	157	GLN
1	A	179	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	251	LYS
1	A	252	ILE
1	A	286	SER
1	A	296	LYS
1	A	298	SER
1	A	300	LYS
1	A	301	LYS
1	A	318	LEU
1	A	356	MET
1	A	358	ASN
1	A	382	LEU
1	A	398	LEU
1	A	399	VAL
1	A	407	VAL
1	A	426	ARG
1	A	439	ARG
1	A	446	ASP
1	A	448	MET
1	A	451	GLN
1	A	455	THR
1	A	496	LEU
1	A	501	LEU
1	A	508	ARG
1	A	523	LYS
1	A	532	ARG
1	A	560	GLU
1	A	562	SER
1	A	667	SER
1	A	700	LYS
1	A	704	VAL
1	A	706	ARG
1	A	718	ASN
1	A	761	LYS
1	A	772	ARG
1	A	776	LEU
1	A	777	SER
1	A	789	VAL
1	A	794	LEU
1	A	797	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	145	ASN
1	A	157	GLN
1	A	172	GLN
1	A	291	GLN
1	A	299	ASN
1	A	336	ASN
1	A	376	ASN
1	A	428	HIS
1	A	444	ASN
1	A	476	GLN
1	A	486	ASN
1	A	492	ASN
1	A	494	HIS
1	A	503	GLN
1	A	549	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	752/852 (88%)	0.14	37 (4%) 29 28	28, 46, 72, 94	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	THR	6.2
1	A	235	LEU	6.1
1	A	92	VAL	5.5
1	A	233	LEU	5.3
1	A	88	ILE	5.2
1	A	86	ARG	5.2
1	A	87	GLN	5.0
1	A	237	GLY	4.9
1	A	234	VAL	4.5
1	A	27	ASP	4.5
1	A	85	MET	4.4
1	A	28	VAL	4.2
1	A	614	ILE	4.1
1	A	248	TYR	3.4
1	A	93	LEU	3.4
1	A	245	VAL	3.1
1	A	83	ALA	2.9
1	A	84	TRP	2.9
1	A	246	GLU	2.9
1	A	707	PRO	2.9
1	A	762	PRO	2.9
1	A	603	VAL	2.8
1	A	705	ASN	2.7
1	A	789	VAL	2.6
1	A	244	GLU	2.6
1	A	790	ALA	2.6
1	A	724	ARG	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	612	VAL	2.5
1	A	656	GLY	2.4
1	A	301	LYS	2.3
1	A	766	ASP	2.3
1	A	223	VAL	2.2
1	A	658	PRO	2.1
1	A	119	GLU	2.0
1	A	76	ASP	2.0
1	A	227	ARG	2.0
1	A	723	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.