



Full wwPDB X-ray Structure Validation Report i

Sep 3, 2023 – 12:47 PM EDT

PDB ID : 3RQO

Title : Structure of the endothelial nitric oxide synthase heme domain in complex with 6-(((3R,4R)-4-(2-((1S,2R/1R,2S)-2-(3-Chlorophenyl)cyclopropylamino)ethyl)oxy)pyrrolidin-3-yl)methyl)-4-methylpyridin-2-amine

Authors : Li, H.; Delker, S.L.; Poulos, T.L.

Deposited on : 2011-04-28

Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 7032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	404	3228	2053	568	591	16	0	3	0
1	B	402	3198	2034	562	586	16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29473
B	100	ARG	CYS	SEE REMARK 999	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O		0	0
			17	9	5	3			


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O		0	0
			17	9	5	3			

- Molecule 4 is 6-{|[(3R,4R)-4-(2-{|[(1R,2S)-2-(3-chlorophenyl)cyclopropyl]amino}ethoxy)pyrrolidin-3-yl]methyl}-4-methylpyridin-2-amine (three-letter code: X2D) (formula: C₂₂H₂₉ClN₄O).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total		C	Cl	N	O	
			28		22	1	4	1	0
4	B	1	Total		C	Cl	N	O	
			28		22	1	4	1	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0

- Molecule 7 is CACODYLIC ACID (three-letter code: CAD) (formula: C₂H₇AsO₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total As C 3 1 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total As C 3 1 2	0	0

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Zn 1 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	198	Total O 198 198	0	0
9	B	205	Total O 205 205	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, endothelial



- Molecule 1: Nitric oxide synthase, endothelial



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.08 Å 106.64 Å 156.52 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.11 – 2.08 35.11 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.3 (35.11-2.08) 99.3 (35.11-2.08)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ ¹	2.68 (at 2.08 Å)	Xtriage
Refinement program	REFMAC	Depositor
R , R_{free}	0.174 , 0.221 0.189 , 0.234	Depositor DCC
R_{free} test set	2930 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.1	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7032	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, GOL, X2D, CAD, ACT, H4B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.70	0/3327	0.75	1/4531 (0.0%)
1	B	0.73	0/3287	0.75	5/4477 (0.1%)
All	All	0.72	0/6614	0.75	6/9008 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH1	8.18	124.39	120.30
1	B	240	ARG	NE-CZ-NH2	-8.00	116.30	120.30
1	B	189	ARG	NE-CZ-NH1	-6.62	116.99	120.30
1	B	378	LEU	CA-CB-CG	6.15	129.45	115.30
1	A	328	LEU	CA-CB-CG	5.79	128.61	115.30
1	B	189	ARG	NE-CZ-NH2	5.70	123.15	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3228	0	3140	30	0
1	B	3198	0	3104	42	0
2	A	43	0	30	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	43	0	30	2	0
3	A	17	0	15	1	0
3	B	17	0	15	0	0
4	A	28	0	29	2	0
4	B	28	0	29	2	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	6	0	8	0	0
6	B	6	0	8	0	0
7	A	3	0	0	3	0
7	B	3	0	0	0	0
8	A	1	0	0	0	0
9	A	198	0	0	2	0
9	B	205	0	0	7	0
All	All	7032	0	6414	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CYS:SG	7:A:950:CAD:AS	2.59	1.21
1:B:236:ARG:HD3	1:B:242:ASP:OD1	1.62	1.00
1:B:236:ARG:HD3	1:B:242:ASP:CG	2.12	0.68
1:A:340:ASN:H	1:A:340:ASN:HD22	1.41	0.67
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.76	0.66
1:B:363:GLU:OE1	4:B:800:X2D:H14	1.95	0.66
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.37	0.64
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.38	0.63
1:A:388:ASP:OD1	1:A:390:ARG:HG3	2.01	0.61
1:B:240:ARG:HD2	1:B:241:GLY:O	2.00	0.61
1:A:384:CYS:CB	7:A:950:CAD:AS	3.09	0.60
1:B:126:LEU:HD11	1:B:156:GLU:HG2	1.83	0.60
1:B:330:LEU:HB2	9:B:1008:HOH:O	2.02	0.60
1:B:236:ARG:CD	1:B:242:ASP:OD1	2.43	0.60
1:B:240:ARG:NH2	9:B:1199:HOH:O	2.31	0.59
1:B:379:GLU:HB2	9:B:1136:HOH:O	2.03	0.58
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.85	0.57
1:A:96:CYS:HB3	1:B:96:CYS:HB3	1.86	0.55
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.35	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:LEU:HB2	9:A:1010:HOH:O	2.06	0.55
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	1.90	0.54
1:A:95:PRO:HB3	1:A:108:PRO:HB2	1.90	0.53
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.90	0.53
1:B:362:THR:HA	1:B:405:ASN:HD21	1.74	0.53
1:B:378:LEU:HB2	9:B:1016:HOH:O	2.07	0.52
1:A:143:SER:HB2	9:A:1018:HOH:O	2.09	0.52
1:B:308:PRO:HD2	1:B:311:LEU:HD12	1.91	0.52
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.93	0.51
1:B:362:THR:HA	1:B:405:ASN:ND2	2.26	0.51
1:B:95:PRO:HG3	1:B:108:PRO:HG2	1.92	0.50
1:A:69:LYS:HA	1:A:69:LYS:HE2	1.95	0.49
1:A:465:GLU:HB3	1:B:105:LEU:HD22	1.95	0.49
1:A:384:CYS:HB3	7:A:950:CAD:AS	2.73	0.48
1:B:99:ARG:HG2	1:B:100:ARG:HD2	1.96	0.48
1:A:249:GLN:HB2	1:A:252:ARG:HG3	1.95	0.48
1:B:249:GLN:HB2	1:B:252:ARG:HG3	1.97	0.47
1:A:340:ASN:O	1:A:478:GLN:HB2	2.15	0.46
1:B:457:SER:HA	1:B:462:PHE:CG	2.50	0.46
1:A:205:SER:O	1:A:209:MET:HG3	2.16	0.46
1:A:264:ARG:HD3	1:A:285:ASN:O	2.15	0.46
1:B:476:ARG:NH1	9:B:1071:HOH:O	2.48	0.46
1:B:231:THR:O	1:B:353:ALA:HA	2.15	0.45
1:A:363:GLU:OE1	4:A:800:X2D:H14	2.17	0.45
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.47	0.45
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.99	0.45
1:B:290:VAL:HG23	9:B:1096:HOH:O	2.18	0.44
1:B:381:VAL:HG21	1:B:404:ILE:HD11	2.00	0.44
1:B:323:GLU:CD	1:B:323:GLU:H	2.21	0.43
1:B:366:THR:O	1:B:370:CYS:HB2	2.19	0.42
1:B:319:HIS:CG	1:B:320:PRO:HD2	2.54	0.42
1:B:308:PRO:HA	1:B:309:PRO:HD3	1.94	0.42
1:B:480:ASP:HA	1:B:481:PRO:HD3	1.91	0.42
1:A:358:TRP:O	4:A:800:X2D:H25	2.20	0.42
1:A:480:ASP:HA	1:A:481:PRO:HD3	1.89	0.42
1:B:359:TYR:CD2	1:B:364:ILE:HD11	2.54	0.42
1:A:105:LEU:HD22	1:B:465:GLU:HB3	2.01	0.42
1:B:457:SER:HA	1:B:462:PHE:CB	2.50	0.42
1:A:449:TRP:HA	3:A:600:H4B:N1	2.35	0.41
2:A:500:HEM:HBC2	2:A:500:HEM:CMC	2.50	0.41
1:B:121:PRO:HG2	1:B:123:GLU:OE2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:GLU:OE1	1:B:472:SER:OG	2.32	0.41
1:B:244:ARG:HD2	1:B:351:SER:HB2	2.02	0.41
2:A:500:HEM:HHC	2:A:500:HEM:CBB	2.50	0.41
1:B:455:SER:O	1:B:456:GLY:C	2.59	0.41
1:A:340:ASN:HD22	1:A:340:ASN:N	2.07	0.41
1:A:392:THR:HB	1:B:423:HIS:HB2	2.03	0.41
1:B:183:ALA:HA	1:B:184:PRO:HD3	1.85	0.41
1:B:340:ASN:O	1:B:478:GLN:HG2	2.19	0.41
4:B:800:X2D:H26	9:B:1205:HOH:O	2.20	0.41
1:B:340:ASN:HD22	1:B:340:ASN:H	1.67	0.41
1:A:126:LEU:HD12	1:A:126:LEU:HA	1.86	0.41
1:A:158:GLU:OE1	1:A:166:HIS:HD2	2.04	0.41
1:A:246:TRP:CD1	1:A:481:PRO:HG2	2.56	0.41
1:B:185:ARG:HB3	1:B:449:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	403/443 (91%)	393 (98%)	10 (2%)	0	100 100
1	B	398/443 (90%)	384 (96%)	14 (4%)	0	100 100
All	All	801/886 (90%)	777 (97%)	24 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	346/376 (92%)	331 (96%)	15 (4%)	29 28
1	B	342/376 (91%)	321 (94%)	21 (6%)	18 15
All	All	688/752 (92%)	652 (95%)	36 (5%)	22 20

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	100	ARG
1	A	102	LEU
1	A	123	GLU
1	A	125	LEU
1	A	126	LEU
1	A	143	SER
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	331	ARG
1	A	340	ASN
1	A	390	ARG
1	A	416	LYS
1	A	458	LEU
1	B	69	LYS
1	B	99	ARG
1	B	100	ARG
1	B	102	LEU
1	B	105	LEU
1	B	126	LEU
1	B	141	LYS
1	B	149	GLU
1	B	240	ARG
1	B	260	ASP
1	B	278	GLN
1	B	323	GLU
1	B	330	LEU
1	B	331	ARG
1	B	340	ASN
1	B	372	PRO
1	B	378	LEU
1	B	379	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	389	THR
1	B	391	THR
1	B	458	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	GLN
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	ACT	B	860	-	3,3,3	0.77	0	3,3,3	0.75	0
6	GOL	B	880	-	5,5,5	0.42	0	5,5,5	0.76	0
2	HEM	A	500	1	41,50,50	1.79	6 (14%)	45,82,82	1.36	8 (17%)
4	X2D	A	800	-	29,31,31	1.23	2 (6%)	33,43,43	2.34	10 (30%)
3	H4B	A	600	-	16,18,18	1.05	1 (6%)	11,26,26	2.80	7 (63%)
4	X2D	B	800	-	29,31,31	0.88	2 (6%)	33,43,43	2.22	9 (27%)
6	GOL	A	880	-	5,5,5	0.47	0	5,5,5	0.15	0
7	CAD	A	950	-	0,2,4	-	-	0,1,6	-	-
7	CAD	B	950	-	0,2,4	-	-	0,1,6	-	-
5	ACT	A	860	-	3,3,3	0.92	0	3,3,3	0.49	0
2	HEM	B	500	1	41,50,50	1.93	7 (17%)	45,82,82	1.87	8 (17%)
3	H4B	B	600	-	16,18,18	1.07	1 (6%)	11,26,26	2.48	6 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	880	-	-	1/4/4/4	-
2	HEM	A	500	1	-	0/12/54/54	-
4	X2D	A	800	-	-	2/13/30/30	0/4/4/4
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	X2D	B	800	-	-	3/13/30/30	0/4/4/4
6	GOL	A	880	-	-	2/4/4/4	-
2	HEM	B	500	1	-	2/12/54/54	-
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3D-C2D	7.68	1.53	1.36
2	A	500	HEM	C3D-C2D	6.76	1.51	1.36
2	B	500	HEM	C3C-C2C	-3.68	1.35	1.40
4	A	800	X2D	C15-C13	3.49	1.54	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3C-CAC	3.38	1.54	1.47
2	A	500	HEM	C3C-C2C	-3.36	1.35	1.40
2	A	500	HEM	C3C-CAC	3.30	1.54	1.47
2	B	500	HEM	CAB-C3B	2.91	1.55	1.47
2	A	500	HEM	CAB-C3B	2.90	1.55	1.47
2	B	500	HEM	FE-ND	2.86	2.11	1.96
3	A	600	H4B	C7-C6	2.55	1.54	1.52
3	B	600	H4B	C7-C6	2.50	1.54	1.52
2	A	500	HEM	CMD-C2D	2.44	1.56	1.50
4	A	800	X2D	C23-CL23	2.42	1.79	1.74
2	A	500	HEM	CMB-C2B	2.34	1.55	1.50
2	B	500	HEM	CMB-C2B	2.26	1.55	1.50
2	B	500	HEM	FE-NB	2.13	2.07	1.96
4	B	800	X2D	C23-CL23	2.10	1.79	1.74
4	B	800	X2D	C15-C13	2.07	1.52	1.49

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	X2D	C15-C14-C21	-6.66	109.43	122.24
4	B	800	X2D	C15-C14-C21	-6.53	109.68	122.24
4	A	800	X2D	C02-N01-C06	6.33	122.90	118.10
2	B	500	HEM	CBA-CAA-C2A	-5.45	103.32	112.62
2	B	500	HEM	C4D-ND-C1D	5.17	110.41	105.07
4	B	800	X2D	C15-C13-N12	-4.93	107.89	117.50
3	A	600	H4B	C8A-C4A-C4	4.66	118.71	114.57
4	B	800	X2D	C02-N01-C06	4.65	121.62	118.10
2	B	500	HEM	C4B-CHC-C1C	4.22	128.13	122.56
4	B	800	X2D	C5'-N1'-C2'	4.08	115.05	105.42
3	A	600	H4B	C4-C4A-N5	3.96	122.44	119.12
3	B	600	H4B	C8A-C4A-C4	3.89	118.03	114.57
4	A	800	X2D	C11-N12-C13	3.85	119.65	114.20
3	B	600	H4B	C4-C4A-N5	3.84	122.34	119.12
2	A	500	HEM	C4D-ND-C1D	3.83	109.03	105.07
4	A	800	X2D	C04-C05-C06	-3.70	117.90	120.32
4	A	800	X2D	C15-C13-N12	-3.66	110.37	117.50
2	B	500	HEM	CBD-CAD-C3D	-3.50	102.90	112.63
3	A	600	H4B	N1-C2-N3	-3.47	119.98	125.42
3	A	600	H4B	C2-N3-C4	3.44	121.39	115.93
2	A	500	HEM	CBA-CAA-C2A	-3.42	106.78	112.62
3	B	600	H4B	N1-C2-N3	-3.25	120.32	125.42
4	A	800	X2D	C5'-N1'-C2'	3.10	112.73	105.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	X2D	C06-C08-C3'	-3.06	105.48	115.55
3	B	600	H4B	C2-N1-C8A	2.97	121.19	114.54
3	A	600	H4B	C2-N1-C8A	2.93	121.11	114.54
2	B	500	HEM	C1B-NB-C4B	2.81	107.98	105.07
3	B	600	H4B	N2-C2-N3	2.76	121.54	117.25
4	B	800	X2D	C04-C05-C06	-2.70	118.55	120.32
2	A	500	HEM	C4B-CHC-C1C	2.58	125.97	122.56
2	A	500	HEM	C1B-NB-C4B	2.56	107.72	105.07
4	A	800	X2D	C05-C06-N01	-2.43	120.32	122.90
4	A	800	X2D	N02-C02-N01	2.37	120.24	116.49
4	B	800	X2D	C26-C21-C14	-2.33	116.67	121.08
3	B	600	H4B	C2-N3-C4	2.31	119.59	115.93
3	A	600	H4B	C4A-N5-C6	-2.30	114.91	121.16
2	A	500	HEM	CMD-C2D-C1D	2.29	128.52	125.04
4	B	800	X2D	N02-C02-N01	2.29	120.10	116.49
2	A	500	HEM	C1D-C2D-C3D	-2.26	104.58	106.96
2	B	500	HEM	CMD-C2D-C1D	2.23	128.43	125.04
4	B	800	X2D	C05-C06-N01	-2.19	120.58	122.90
4	B	800	X2D	C06-C08-C3'	-2.15	108.47	115.55
3	A	600	H4B	N2-C2-N3	2.11	120.53	117.25
2	A	500	HEM	CHA-C4D-ND	2.09	126.96	124.38
4	A	800	X2D	C05-C04-C03	2.08	120.56	118.09
2	B	500	HEM	CHA-C4D-ND	2.06	126.93	124.38
2	A	500	HEM	CAD-C3D-C4D	2.04	128.23	124.66
2	B	500	HEM	CMA-C3A-C4A	-2.04	125.32	128.46

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	880	GOL	C1-C2-C3-O3
6	A	880	GOL	O1-C1-C2-C3
6	A	880	GOL	O1-C1-C2-O2
4	A	800	X2D	C15-C14-C21-C26
4	A	800	X2D	C15-C14-C21-C22
4	B	800	X2D	O09-C10-C11-N12
4	B	800	X2D	C15-C14-C21-C22
4	B	800	X2D	C15-C14-C21-C26
2	B	500	HEM	CAA-CBA-CGA-O2A
2	B	500	HEM	CAA-CBA-CGA-O1A



There are no ring outliers.


6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	3	0
4	A	800	X2D	2	0
3	A	600	H4B	1	0
4	B	800	X2D	2	0
7	A	950	CAD	3	0
2	B	500	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/443 (91%)	0.39	34 (8%) 11 13	26, 38, 61, 81	0
1	B	402/443 (90%)	0.23	19 (4%) 31 36	25, 39, 61, 83	0
All	All	806/886 (90%)	0.31	53 (6%) 18 22	25, 38, 61, 83	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	5.9
1	A	259	GLN	5.6
1	B	259	GLN	4.8
1	A	123	GLU	4.6
1	A	124	GLN	4.5
1	A	122	ALA	4.3
1	A	160	ALA	4.1
1	A	68	PRO	3.9
1	B	146	GLN	3.9
1	A	69	LYS	3.8
1	A	153	GLN	3.8
1	B	121	PRO	3.7
1	A	261	GLY	3.6
1	A	257	ARG	3.5
1	A	121	PRO	3.4
1	B	239	GLY	3.3
1	B	147	ALA	3.3
1	A	260	ASP	3.1
1	A	230	ILE	3.0
1	B	142	ARG	3.0
1	A	232	VAL	2.9
1	A	337	ALA	2.9
1	B	122	ALA	2.8
1	B	141	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	260	ASP	2.7
1	B	261	GLY	2.6
1	A	127	SER	2.6
1	B	323	GLU	2.6
1	A	91	GLN	2.6
1	A	125	LEU	2.5
1	A	126	LEU	2.5
1	A	67	GLY	2.5
1	B	123	GLU	2.5
1	A	338	VAL	2.4
1	A	390	ARG	2.4
1	B	144	GLY	2.4
1	A	156	GLU	2.4
1	A	231	THR	2.4
1	B	145	SER	2.4
1	A	258	GLN	2.3
1	A	278	GLN	2.2
1	A	364	ILE	2.2
1	B	449	TRP	2.2
1	B	386	ASP	2.2
1	A	107	LEU	2.2
1	B	390	ARG	2.2
1	A	157	ALA	2.1
1	A	213	ILE	2.1
1	A	200	ALA	2.1
1	B	149	GLU	2.1
1	A	335	LEU	2.0
1	A	454	ILE	2.0
1	B	140	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	A	880	6/6	0.85	0.21	55,56,57,57	0
6	GOL	B	880	6/6	0.87	0.20	52,53,55,55	0
4	X2D	A	800	28/28	0.92	0.17	29,38,41,47	0
5	ACT	A	860	4/4	0.94	0.19	45,46,48,48	0
4	X2D	B	800	28/28	0.96	0.16	33,38,47,53	0
5	ACT	B	860	4/4	0.97	0.26	47,49,49,51	0
2	HEM	A	500	43/43	0.98	0.17	21,28,34,39	0
2	HEM	B	500	43/43	0.98	0.16	27,30,35,36	0
3	H4B	A	600	17/17	0.98	0.15	23,28,31,31	0
7	CAD	A	950	3/5	0.98	0.07	70,70,70,71	0
7	CAD	B	950	3/5	0.98	0.22	77,77,77,77	0
3	H4B	B	600	17/17	0.99	0.16	23,28,31,34	0
8	ZN	A	900	1/1	1.00	0.08	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.