



wwPDB X-ray Structure Validation Summary Report

Sep 4, 2023 – 05:18 PM EDT

PDB ID : 3TSW
Title : crystal structure of the PDZ3-SH3-GUK core module of Human ZO-1
Authors : Nomme, J.; Lavie, A.
Deposited on : 2011-09-13
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	
1	B	391	
1	C	391	
1	D	391	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tight junction protein ZO-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	334	Total 2687	C 1709	N 481	O 491	S 6	0	0	0
1	B	194	Total 1552	C 991	N 270	O 290	S 1	0	0	0
1	C	337	Total 2703	C 1714	N 488	O 495	S 6	0	0	0
1	D	199	Total 1590	C 1011	N 281	O 297	S 1	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	413	GLY	-	expression tag	UNP Q07157
A	414	SER	-	expression tag	UNP Q07157
A	415	HIS	-	expression tag	UNP Q07157
A	416	MET	-	expression tag	UNP Q07157
B	413	GLY	-	expression tag	UNP Q07157
B	414	SER	-	expression tag	UNP Q07157
B	415	HIS	-	expression tag	UNP Q07157
B	416	MET	-	expression tag	UNP Q07157
C	413	GLY	-	expression tag	UNP Q07157
C	414	SER	-	expression tag	UNP Q07157
C	415	HIS	-	expression tag	UNP Q07157
C	416	MET	-	expression tag	UNP Q07157
D	413	GLY	-	expression tag	UNP Q07157
D	414	SER	-	expression tag	UNP Q07157
D	415	HIS	-	expression tag	UNP Q07157
D	416	MET	-	expression tag	UNP Q07157

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total	O	0	0
			20	20		
3	B	13	Total	O	0	0
			13	13		
3	C	22	Total	O	0	0
			22	22		
3	D	15	Total	O	0	0
			15	15		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	100.63Å 100.63Å 182.52Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	87.17 – 2.85 29.55 – 2.85	Depositor EDS
% Data completeness (in resolution range)	96.2 (87.17-2.85) 97.6 (29.55-2.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.211 , 0.290 0.191 , 0.251	Depositor DCC
R_{free} test set	4762 reflections (10.10%)	wwPDB-VP
Wilson B-factor (Å ²)	75.9	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.002 for h,-h-k,-l 0.139 for -k,-h,-l	Xtriage
Reported twinning fraction	0.508 for 1.000H, 1.000K, L 0.492 for -1.000H, 1.000H+1.000K, -L	Depositor
Outliers	0 of 47162 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8637	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2736	0.71	0/3688
1	B	0.62	0/1578	0.79	1/2124 (0.0%)
1	C	0.55	0/2753	0.72	2/3712 (0.1%)
1	D	0.68	0/1615	0.85	2/2175 (0.1%)
All	All	0.59	0/8682	0.76	5/11699 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	488	LEU	CA-CB-CG	5.31	127.51	115.30
1	B	796	GLN	N-CA-C	5.25	125.18	111.00
1	C	424	LEU	CA-CB-CG	5.21	127.29	115.30
1	D	489	ASP	CB-CG-OD2	5.09	122.88	118.30
1	C	797	LEU	CA-CB-CG	5.08	126.98	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	482	GLU	Peptide
1	B	795	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2687	0	2722	112	0
1	B	1552	0	1564	90	0
1	C	2703	0	2722	106	0
1	D	1590	0	1605	100	0
2	A	10	0	0	1	0
2	B	10	0	0	2	0
2	C	5	0	0	0	0
2	D	10	0	0	0	0
3	A	20	0	0	4	0
3	B	13	0	0	0	0
3	C	22	0	0	6	0
3	D	15	0	0	3	0
All	All	8637	0	8613	387	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 387 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:450:VAL:HG12	1:D:451:LEU:H	1.19	1.08
1:A:558:LEU:HD12	1:A:573:ILE:HG22	1.28	1.05
1:B:582:GLN:NE2	1:D:478:ILE:HD12	1.77	0.99
1:B:518:SER:OG	2:B:7:SO4:O4	1.88	0.92
1:B:431:ASP:H	1:B:492:LYS:HG2	1.33	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/391 (84%)	266 (81%)	48 (15%)	14 (4%)	2	5
1	B	188/391 (48%)	153 (81%)	29 (15%)	6 (3%)	4	9
1	C	331/391 (85%)	265 (80%)	53 (16%)	13 (4%)	3	6
1	D	193/391 (49%)	150 (78%)	35 (18%)	8 (4%)	3	5
All	All	1040/1564 (66%)	834 (80%)	165 (16%)	41 (4%)	3	6

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	515	VAL
1	A	631	PRO
1	A	689	ILE
1	C	454	SER
1	C	676	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/338 (86%)	248 (86%)	41 (14%)	3	6
1	B	167/338 (49%)	130 (78%)	37 (22%)	1	0
1	C	289/338 (86%)	243 (84%)	46 (16%)	2	4
1	D	171/338 (51%)	144 (84%)	27 (16%)	2	4
All	All	916/1352 (68%)	765 (84%)	151 (16%)	2	3

5 of 151 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	761	LEU
1	D	571	ARG
1	C	770	THR
1	D	458	LYS
1	D	792	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	796	GLN
1	D	793	GLN
1	D	582	GLN
1	B	582	GLN
1	C	793	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	6	-	4,4,4	0.19	0	6,6,6	0.12	0
2	SO4	A	5	-	4,4,4	0.14	0	6,6,6	0.22	0
2	SO4	B	7	-	4,4,4	0.27	0	6,6,6	0.43	0
2	SO4	D	4	-	4,4,4	0.12	0	6,6,6	0.34	0
2	SO4	C	3	-	4,4,4	0.15	0	6,6,6	0.33	0
2	SO4	D	2	-	4,4,4	0.29	0	6,6,6	0.45	0
2	SO4	B	1	-	4,4,4	0.27	0	6,6,6	0.31	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5	SO4	1	0
2	B	7	SO4	1	0
2	B	1	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/391 (85%)	-0.20	3 (0%) 84 83	53, 74, 86, 89	0
1	B	194/391 (49%)	-0.18	2 (1%) 82 79	52, 65, 82, 85	0
1	C	337/391 (86%)	-0.10	5 (1%) 73 70	52, 76, 92, 100	0
1	D	199/391 (50%)	-0.29	0 100 100	47, 62, 85, 91	0
All	All	1064/1564 (68%)	-0.18	10 (0%) 84 83	47, 71, 89, 100	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	572	GLY	3.3
1	B	593	THR	3.1
1	A	718	TYR	2.9
1	C	573	ILE	2.5
1	C	571	ARG	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	A	6	5/5	0.92	0.10	109,110,111,111	0
2	SO4	B	7	5/5	0.95	0.16	70,70,71,71	0
2	SO4	A	5	5/5	0.96	0.08	97,97,98,98	0
2	SO4	B	1	5/5	0.98	0.09	64,65,66,67	0
2	SO4	C	3	5/5	0.98	0.12	67,67,69,70	0
2	SO4	D	2	5/5	0.98	0.11	55,57,60,61	0
2	SO4	D	4	5/5	0.98	0.18	72,73,74,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.