



# wwPDB X-ray Structure Validation Summary Report i

Oct 5, 2023 – 04:44 AM EDT

PDB ID : 6VQO  
Title : T cell receptor-p53-HLA-A2 complex  
Authors : Wu, D.; Gallagher, D.T.; Pierce, B.G.; Mariuzza, R.A.  
Deposited on : 2020-02-05  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	: FAILED
Xtriage (Phenix)	: 1.13
EDS	: FAILED
Percentile statistics	: 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.35.1

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbit and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 12740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C 2198	N 1378	O 395	S 416	9	0	0
1	F	254	Total	C 1988	N 1250	O 355	S 375	8	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP F6IQS2
A	276	GLY	-	expression tag	UNP F6IQS2
A	277	GLY	-	expression tag	UNP F6IQS2
A	278	GLY	-	expression tag	UNP F6IQS2
A	279	LEU	-	expression tag	UNP F6IQS2
A	280	ASN	-	expression tag	UNP F6IQS2
A	281	ASP	-	expression tag	UNP F6IQS2
A	282	ILE	-	expression tag	UNP F6IQS2
A	283	PHE	-	expression tag	UNP F6IQS2
A	284	GLU	-	expression tag	UNP F6IQS2
A	285	ALA	-	expression tag	UNP F6IQS2
A	286	GLN	-	expression tag	UNP F6IQS2
A	287	LYS	-	expression tag	UNP F6IQS2
A	288	ILE	-	expression tag	UNP F6IQS2
A	289	GLU	-	expression tag	UNP F6IQS2
A	290	TRP	-	expression tag	UNP F6IQS2
A	291	HIS	-	expression tag	UNP F6IQS2
A	292	GLU	-	expression tag	UNP F6IQS2
F	0	MET	-	expression tag	UNP F6IQS2
F	276	GLY	-	expression tag	UNP F6IQS2
F	277	GLY	-	expression tag	UNP F6IQS2
F	278	GLY	-	expression tag	UNP F6IQS2
F	279	LEU	-	expression tag	UNP F6IQS2
F	280	ASN	-	expression tag	UNP F6IQS2
F	281	ASP	-	expression tag	UNP F6IQS2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	282	ILE	-	expression tag	UNP F6IQS2
F	283	PHE	-	expression tag	UNP F6IQS2
F	284	GLU	-	expression tag	UNP F6IQS2
F	285	ALA	-	expression tag	UNP F6IQS2
F	286	GLN	-	expression tag	UNP F6IQS2
F	287	LYS	-	expression tag	UNP F6IQS2
F	288	ILE	-	expression tag	UNP F6IQS2
F	289	GLU	-	expression tag	UNP F6IQS2
F	290	TRP	-	expression tag	UNP F6IQS2
F	291	HIS	-	expression tag	UNP F6IQS2
F	292	GLU	-	expression tag	UNP F6IQS2

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			812	520	137	151	4			
2	G	99	Total	C	N	O	S	0	0	0
			785	504	132	146	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P61769
G	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called T-cell receptor 1a2, alfa chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	193	Total	C	N	O	S	0	0	0
			1493	941	237	305	10			
3	H	196	Total	C	N	O	S	0	0	0
			1514	948	244	312	10			

- Molecule 4 is a protein called TCR receptor 1a2, beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	240	Total	C	N	O	S	0	0	0
			1882	1190	325	359	8			
4	J	241	Total	C	N	O	S	0	0	0
			1916	1212	333	363	8			

- Molecule 5 is a protein called peptide from p53 tumor suppressor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	P	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			
5	Q	9	Total	C	N	O	S	0	0	0
			76	45	16	13	2			

MolProbity and EDS failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.13 Å    118.13 Å    153.13 Å 90.00°    90.00°    120.00°	Depositor
Resolution (Å)	48.52 – 3.00	Depositor
% Data completeness (in resolution range)	97.2 (48.52-3.00)	Depositor
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.26 (at 3.01 Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
$R$ , $R_{free}$	0.162 , 0.211	Depositor
Wilson B-factor (Å <sup>2</sup> )	56.4	Xtriage
Anisotropy	0.010	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.39$ , $< L^2 > = 0.22$	Xtriage
Estimated twinning fraction	0.098 for -h,-k,l 0.378 for h,-h-k,-l 0.098 for -k,-h,-l	Xtriage
Total number of atoms	12740	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [\(i\)](#)

### 4.1 Standard geometry [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [\(i\)](#)

#### 4.3.1 Protein backbone [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [\(i\)](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 4.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data [\(i\)](#)

### 5.1 Protein, DNA and RNA chains [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands [\(i\)](#)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers [\(i\)](#)

EDS failed to run properly - this section is therefore empty.