



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 24, 2021 – 06:14 pm GMT

PDB ID : 6Z9N
Title : Copper transporter OprC
Authors : Bhamidimarri, S.P.; van den Berg, B.
Deposited on : 2020-06-04
Resolution : 2.73 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	723	 70% 18% • 10%
1	B	723	 2% 68% 21% • 9%
1	C	723	 2% 63% 23% • 11%
1	D	723	 2% 67% 21% • 9%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 20292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative copper transport outer membrane porin OprC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	648	5028	3156	885	970	17	4	1	0
1	B	657	5092	3194	899	980	19	0	1	0
1	C	644	5001	3141	882	963	15	4	1	0
1	D	658	5100	3200	900	981	19	4	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	323	ALA	HIS	engineered mutation	UNP G3XD89
B	323	ALA	HIS	engineered mutation	UNP G3XD89
C	323	ALA	HIS	engineered mutation	UNP G3XD89
D	323	ALA	HIS	engineered mutation	UNP G3XD89

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Cu 1	0	0
2	B	1	Total 1	Cu 1	0	0
2	C	1	Total 1	Cu 1	0	0
2	D	1	Total 1	Cu 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total 28	O 28	0	0
3	B	13	Total 13	O 13	0	0
3	C	17	Total 17	O 17	0	0
3	D	9	Total 9	O 9	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative copper transport outer membrane porin OprC



- Molecule 1: Putative copper transport outer membrane porin OprC





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.33Å 197.27Å 172.12Å 90.00° 91.42° 90.00°	Depositor
Resolution (Å)	67.30 – 2.73 172.07 – 2.73	Depositor EDS
% Data completeness (in resolution range)	98.3 (67.30-2.73) 92.9 (172.07-2.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.73Å)	Xtrriage
Refinement program	PHENIX 1.18_3855	Depositor
R, R_{free}	0.216 , 0.267 0.216 , 0.267	Depositor DCC
R_{free} test set	5947 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtrriage
Anisotropy	0.424	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.226 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	20292	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	1/5155 (0.0%)	0.71	1/6990 (0.0%)
1	B	0.50	0/5222	0.67	1/7079 (0.0%)
1	C	0.51	0/5123	0.69	2/6941 (0.0%)
1	D	0.48	0/5230	0.68	1/7090 (0.0%)
All	All	0.50	1/20730 (0.0%)	0.69	5/28100 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	375	ARG	CG-CD	5.06	1.64	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	147	MET	CB-CG-SD	7.23	134.09	112.40
1	B	147	MET	CB-CG-SD	6.18	130.93	112.40
1	D	147	MET	CG-SD-CE	5.41	108.86	100.20
1	C	147	MET	CG-SD-CE	5.08	108.33	100.20
1	A	348	ASP	CB-CG-OD1	5.00	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5028	0	4808	71	1
1	B	5092	0	4870	89	1
1	C	5001	0	4791	102	0
1	D	5100	0	4881	93	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	28	0	0	2	0
3	B	13	0	0	0	0
3	C	17	0	0	1	0
3	D	9	0	0	1	0
All	All	20292	0	19350	347	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 347 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:TYR:OH	1:A:390:ASP:OD1	1.88	0.91
1:A:509:SER:HB3	1:A:515:LYS:HG3	1.62	0.81
1:B:202:LEU:HD12	1:C:225:LEU:HD22	1.63	0.80
1:C:94:PRO:O	1:C:209:ARG:NH2	2.13	0.80
1:A:672:ARG:H	1:A:672:ARG:HD2	1.49	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:713:ARG:NH1	1:B:623:GLU:OE2[2_646]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	645/723 (89%)	601 (93%)	34 (5%)	10 (2%)	9	17
1	B	656/723 (91%)	624 (95%)	27 (4%)	5 (1%)	19	36
1	C	633/723 (88%)	578 (91%)	47 (7%)	8 (1%)	12	21
1	D	657/723 (91%)	605 (92%)	44 (7%)	8 (1%)	13	24
All	All	2591/2892 (90%)	2408 (93%)	152 (6%)	31 (1%)	13	24

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	478	ASP
1	A	555	ARG
1	A	625	ASP
1	C	336	SER
1	C	389	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	519/572 (91%)	485 (93%)	34 (7%)	16	29
1	B	525/572 (92%)	487 (93%)	38 (7%)	14	25
1	C	517/572 (90%)	473 (92%)	44 (8%)	10	20
1	D	526/572 (92%)	483 (92%)	43 (8%)	11	21
All	All	2087/2288 (91%)	1928 (92%)	159 (8%)	13	23

5 of 159 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	721	PHE
1	D	510	VAL
1	D	139	MET
1	D	307	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	586	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	76	GLN
1	D	408	ASN
1	D	705	ASN
1	D	547	GLN
1	B	199	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	648/723 (89%)	0.09	2 (0%) 94 96	43, 56, 86, 125	0
1	B	657/723 (90%)	0.24	10 (1%) 73 79	42, 60, 96, 124	0
1	C	644/723 (89%)	0.07	5 (0%) 86 89	42, 61, 100, 130	0
1	D	658/723 (91%)	0.25	15 (2%) 60 67	43, 62, 100, 135	0
All	All	2607/2892 (90%)	0.16	32 (1%) 79 83	42, 60, 96, 135	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	362	TRP	5.3
1	C	510	VAL	4.2
1	A	555	ARG	4.1
1	D	362	TRP	3.9
1	B	387	MET	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CU	D	801	1/1	0.96	0.14	97,97,97,97	0
2	CU	A	801	1/1	0.97	0.16	83,83,83,83	0
2	CU	B	801	1/1	0.98	0.13	88,88,88,88	0
2	CU	C	801	1/1	0.99	0.14	95,95,95,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.