



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 09:26 PM EDT

PDB ID : 5ZL5
Title : Crystal structure of DFA-IIIase mutant C387A from *Arthrobacter chlorophenolicus* A6
Authors : Yu, S.H.; Shen, H.; Li, X.; Mu, W.M.
Deposited on : 2018-03-26
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<div><div>2%</div><div>94%</div><div>5%</div><div>• •</div></div>
1	B	445	<div><div>%</div><div>94%</div><div>5%</div><div>• •</div></div>
1	C	445	<div><div>3%</div><div>93%</div><div>5%</div><div>•</div></div>
1	D	445	<div><div>%</div><div>94%</div><div>5%</div><div>• •</div></div>
1	E	445	<div><div>2%</div><div>92%</div><div>6%</div><div>• •</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	445	<div><div></div><div>2%</div><div>94%</div><div></div><div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 21845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DFA-IIIase C387A mutant.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	0	0
			3281	2042	585	648	6			
1	B	438	Total	C	N	O	S	0	0	0
			3282	2041	585	650	6			
1	C	437	Total	C	N	O	S	0	0	0
			3277	2040	584	647	6			
1	D	438	Total	C	N	O	S	0	1	0
			3286	2047	585	648	6			
1	E	436	Total	C	N	O	S	0	0	0
			3272	2036	583	647	6			
1	F	435	Total	C	N	O	S	0	1	0
			3267	2034	582	645	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	387	ALA	CYS	engineered mutation	UNP B8HDZ1
B	387	ALA	CYS	engineered mutation	UNP B8HDZ1
C	387	ALA	CYS	engineered mutation	UNP B8HDZ1
D	387	ALA	CYS	engineered mutation	UNP B8HDZ1
E	387	ALA	CYS	engineered mutation	UNP B8HDZ1
F	387	ALA	CYS	engineered mutation	UNP B8HDZ1

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	E	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	401	Total	O	0	0
			401	401		
3	B	386	Total	O	0	0
			386	386		
3	C	350	Total	O	0	0
			350	350		
3	D	346	Total	O	0	0
			346	346		
3	E	362	Total	O	0	0
			362	362		
3	F	305	Total	O	0	0
			305	305		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DFA-IIIase C387A mutant



- Molecule 1: DFA-IIIase C387A mutant



- Molecule 1: DFA-IIIase C387A mutant




- Molecule 1: DFA-IIIase C387A mutant





- Molecule 1: DFA-IIIase C387A mutant





● Molecule 1: DFA-IIIase C387A mutant



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	116.30Å 79.53Å 140.97Å 90.00° 100.54° 90.00°	Depositor
Resolution (Å)	50.00 – 1.65 46.42 – 1.65	Depositor EDS
% Data completeness (in resolution range)	96.0 (50.00-1.65) 96.0 (46.42-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.132 , 0.173 0.133 , 0.174	Depositor DCC
R_{free} test set	14338 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	12.4	Xtriage
Anisotropy	0.278	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	21845	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.50	0/3353	0.76	1/4572 (0.0%)
1	B	0.51	0/3353	0.78	2/4572 (0.0%)
1	C	0.49	0/3349	0.74	1/4567 (0.0%)
1	D	0.49	0/3361	0.73	1/4582 (0.0%)
1	E	0.48	0/3343	0.74	1/4559 (0.0%)
1	F	0.47	0/3341	0.72	0/4556
All	All	0.49	0/20100	0.74	6/27408 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	61	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	C	6	ARG	NE-CZ-NH2	5.37	122.99	120.30
1	D	164	ALA	CB-CA-C	-5.12	102.41	110.10
1	E	106	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	A	277	ARG	NE-CZ-NH2	-5.05	117.77	120.30
1	B	357	VAL	CG1-CB-CG2	5.01	118.91	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3281	0	3136	6	0
1	B	3282	0	3132	9	0
1	C	3277	0	3133	14	0
1	D	3286	0	3140	11	0
1	E	3272	0	3124	11	0
1	F	3267	0	3125	8	0
2	A	6	0	8	0	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
2	E	6	0	8	1	0
2	F	6	0	8	0	0
3	A	401	0	0	3	0
3	B	386	0	0	1	0
3	C	350	0	0	3	0
3	D	346	0	0	3	0
3	E	362	0	0	2	0
3	F	305	0	0	1	0
All	All	21845	0	18830	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:275:ASN:HB2	3:D:601:HOH:O	1.68	0.92
1:C:312:PHE:C	3:C:601:HOH:O	2.09	0.90
1:F:275:ASN:HB2	3:F:601:HOH:O	1.71	0.89
1:A:204:HIS:NE2	3:A:601:HOH:O	2.18	0.75
3:E:601:HOH:O	1:F:204:HIS:NE2	2.19	0.74
3:A:601:HOH:O	1:B:204:HIS:NE2	2.20	0.74
3:A:601:HOH:O	1:D:204:HIS:NE2	2.19	0.74
1:C:307:GLU:HG2	1:C:312:PHE:O	1.88	0.73
1:C:307:GLU:CG	1:C:312:PHE:O	2.43	0.67
1:C:309:TRP:O	1:C:312:PHE:O	2.15	0.62
1:C:313:PHE:N	3:C:601:HOH:O	2.34	0.55
1:C:312:PHE:CA	3:C:601:HOH:O	2.54	0.55
1:E:168:THR:HA	1:E:191:ASN:O	2.08	0.53
1:F:168:THR:HA	1:F:191:ASN:O	2.09	0.53
1:E:155:SER:HB3	3:E:707:HOH:O	2.09	0.51
1:C:322:LEU:HA	1:C:357:VAL:HG13	1.91	0.51
1:D:322:LEU:HA	1:D:357:VAL:HG13	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:THR:HA	1:A:191:ASN:O	2.10	0.51
1:B:168:THR:HA	1:B:191:ASN:O	2.11	0.50
1:D:282:TYR:CE1	1:D:316:ASP:HA	2.47	0.49
1:A:85:ILE:HG12	1:A:387:ALA:HB1	1.93	0.49
1:C:282:TYR:CE1	1:C:316:ASP:HA	2.47	0.49
1:C:168:THR:HA	1:C:191:ASN:O	2.12	0.49
1:F:282:TYR:CE1	1:F:316:ASP:HA	2.47	0.49
1:B:322:LEU:HA	1:B:357:VAL:HG13	1.94	0.49
1:D:168:THR:HA	1:D:191:ASN:O	2.12	0.49
1:E:313:PHE:CD1	1:E:383:SER:HB3	2.48	0.48
1:C:220:TRP:CD1	2:E:501:GOL:H11	2.48	0.48
1:C:205:HIS:HA	1:C:228:ASP:O	2.14	0.47
1:E:322:LEU:HA	1:E:357:VAL:HG13	1.96	0.47
1:E:306:HIS:HD2	1:E:307:GLU:O	1.96	0.47
1:B:282:TYR:CE1	1:B:316:ASP:HA	2.50	0.47
1:E:153:ASP:OD2	1:E:236:ARG:NH2	2.50	0.45
1:C:6:ARG:HD2	1:E:52:TYR:HE2	1.82	0.45
1:D:142:ASN:HA	1:D:183:ASP:O	2.17	0.45
1:F:322:LEU:HA	1:F:357:VAL:HG13	1.99	0.45
1:B:205:HIS:HA	1:B:228:ASP:O	2.17	0.44
1:E:251:THR:OG1	1:F:275:ASN:ND2	2.37	0.43
1:C:274:THR:HA	1:C:298:ALA:O	2.18	0.43
1:D:228:ASP:HA	1:D:252:ALA:O	2.18	0.43
1:A:282:TYR:CE1	1:A:316:ASP:HA	2.54	0.43
1:A:205:HIS:HA	1:A:228:ASP:O	2.19	0.43
1:D:291:ASN:HB2	3:D:887:HOH:O	2.19	0.43
1:B:384:SER:HB3	1:B:390:ALA:HB2	2.01	0.42
1:D:205:HIS:HA	1:D:228:ASP:O	2.19	0.42
1:E:205:HIS:HA	1:E:228:ASP:O	2.20	0.42
1:A:227:THR:HA	1:A:251:THR:O	2.20	0.41
1:F:205:HIS:HA	1:F:228:ASP:O	2.19	0.41
1:B:154:GLY:N	3:B:501:HOH:O	2.52	0.41
1:B:379:VAL:HG12	1:B:400:ALA:HB2	2.03	0.41
1:E:221:GLY:HA3	1:E:245:HIS:CE1	2.56	0.41
1:B:251:THR:OG1	1:D:275:ASN:ND2	2.39	0.41
1:D:156:GLY:HA2	3:D:861:HOH:O	2.21	0.41
1:E:142:ASN:HA	1:E:183:ASP:O	2.21	0.40
1:C:230:LEU:HD23	1:C:254:ASN:HB3	2.03	0.40
1:F:388:PHE:HA	1:F:391:GLN:HE21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/445 (98%)	414 (95%)	20 (5%)	0	100	100
1	B	434/445 (98%)	415 (96%)	19 (4%)	0	100	100
1	C	433/445 (97%)	414 (96%)	19 (4%)	0	100	100
1	D	435/445 (98%)	419 (96%)	16 (4%)	0	100	100
1	E	432/445 (97%)	413 (96%)	19 (4%)	0	100	100
1	F	432/445 (97%)	415 (96%)	17 (4%)	0	100	100
All	All	2600/2670 (97%)	2490 (96%)	110 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/352 (99%)	342 (98%)	7 (2%)	55	32
1	B	349/352 (99%)	343 (98%)	6 (2%)	60	39
1	C	349/352 (99%)	342 (98%)	7 (2%)	55	32
1	D	349/352 (99%)	343 (98%)	6 (2%)	60	39
1	E	348/352 (99%)	340 (98%)	8 (2%)	50	25
1	F	348/352 (99%)	340 (98%)	8 (2%)	50	25
All	All	2092/2112 (99%)	2050 (98%)	42 (2%)	55	32

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	19	LYS
1	A	228	ASP
1	A	304	ARG
1	A	325	LEU
1	A	386	SER
1	A	425	ASP
1	B	4	ASN
1	B	100	LEU
1	B	228	ASP
1	B	304	ARG
1	B	325	LEU
1	B	402	ASP
1	C	42	ASP
1	C	100	LEU
1	C	228	ASP
1	C	304	ARG
1	C	325	LEU
1	C	352	GLU
1	C	402	ASP
1	D	100	LEU
1	D	228	ASP
1	D	291	ASN
1	D	304	ARG
1	D	325	LEU
1	D	350	ARG
1	E	4	ASN
1	E	100	LEU
1	E	118	ASP
1	E	228	ASP
1	E	291	ASN
1	E	304	ARG
1	E	325	LEU
1	E	402	ASP
1	F	100	LEU
1	F	228	ASP
1	F	304	ARG
1	F	325	LEU
1	F	347	ASN
1	F	357	VAL
1	F	375	VAL
1	F	402	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	176	ASN
1	B	4	ASN
1	B	176	ASN
1	C	176	ASN
1	C	291	ASN
1	D	176	ASN
1	D	291	ASN
1	E	4	ASN
1	E	176	ASN
1	E	291	ASN
1	E	306	HIS
1	F	176	ASN
1	F	291	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	C	501	-	5,5,5	0.42	0	5,5,5	0.66	0
2	GOL	E	501	-	5,5,5	0.21	0	5,5,5	0.62	0
2	GOL	A	501	-	5,5,5	0.37	0	5,5,5	0.34	0
2	GOL	F	501	-	5,5,5	0.33	0	5,5,5	0.48	0
2	GOL	D	501	-	5,5,5	0.33	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	501	-	-	0/4/4/4	-
2	GOL	E	501	-	-	0/4/4/4	-
2	GOL	A	501	-	-	0/4/4/4	-
2	GOL	F	501	-	-	0/4/4/4	-
2	GOL	D	501	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/445 (98%)	-0.57	7 (1%) 72 75	5, 11, 27, 71	0
1	B	438/445 (98%)	-0.59	5 (1%) 80 83	6, 11, 28, 61	0
1	C	437/445 (98%)	-0.42	13 (2%) 50 51	7, 12, 33, 73	0
1	D	438/445 (98%)	-0.40	4 (0%) 84 86	7, 13, 31, 60	0
1	E	436/445 (97%)	-0.48	10 (2%) 60 61	8, 14, 33, 45	0
1	F	435/445 (97%)	-0.33	10 (2%) 60 61	9, 15, 37, 68	0
All	All	2622/2670 (98%)	-0.47	49 (1%) 66 69	5, 13, 31, 73	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	156	GLY	5.8
1	F	399	GLU	5.1
1	D	399	GLU	4.8
1	C	400	ALA	4.5
1	E	40	ASP	4.4
1	C	399	GLU	4.3
1	B	399	GLU	4.3
1	A	400	ALA	4.1
1	A	401	ALA	4.1
1	C	113	ALA	4.1
1	F	390	ALA	3.7
1	B	113	ALA	3.6
1	A	399	GLU	3.3
1	F	400	ALA	3.3
1	C	382	ALA	3.3
1	D	156	GLY	3.3
1	F	119	SER	3.0
1	A	113	ALA	3.0
1	E	347	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	401	ALA	2.8
1	F	14	ALA	2.6
1	C	155	SER	2.5
1	B	118	ASP	2.5
1	A	347	ASN	2.4
1	D	398	THR	2.4
1	F	154	GLY	2.4
1	A	443	VAL	2.4
1	C	402	ASP	2.4
1	F	389	GLU	2.3
1	E	39	ALA	2.3
1	C	347	ASN	2.3
1	E	352	GLU	2.3
1	C	156	GLY	2.3
1	B	398	THR	2.3
1	C	386	SER	2.3
1	E	155	SER	2.3
1	E	38	ALA	2.2
1	C	394	ALA	2.2
1	E	41	VAL	2.2
1	F	155	SER	2.2
1	A	442	THR	2.2
1	D	382	ALA	2.2
1	C	387	ALA	2.1
1	B	443	VAL	2.1
1	E	156	GLY	2.1
1	E	118	ASP	2.1
1	E	113	ALA	2.1
1	C	383	SER	2.1
1	F	382	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	C	501	6/6	0.76	0.33	27,28,37,42	0
2	GOL	E	501	6/6	0.85	0.27	26,28,34,35	0
2	GOL	F	501	6/6	0.88	0.22	26,27,33,34	0
2	GOL	A	501	6/6	0.90	0.10	27,28,32,33	0
2	GOL	D	501	6/6	0.94	0.17	26,27,32,33	0

6.5 Other polymers

There are no such residues in this entry.